
Bat Influenza A(HL18NL11) Virus in Fruit Bats, Brazil

Angélica Cristine Almeida Campos,
Luiz Gustavo Bentim Góes, Andres Moreira-Soto,
Cristiano de Carvalho, Guilherme Ambar,
Anna-Lena Sander, Carlo Fischer,
Adriana Ruckert da Rosa,
Debora Cardoso de Oliveira,
Ana Paula G. Kataoka, Wagner André Pedro,
Luzia Fátima A. Martorelli,
Luzia Helena Queiroz, Ariovaldo P. Cruz-Neto,
Edison Luiz Durigon,¹ Jan Felix Drexler¹

Screening of 533 bats for influenza A viruses showed subtype HL18NL11 in intestines of 2 great fruit-eating bats (*Artibeus lituratus*). High concentrations suggested fecal shedding. Genomic characterizations revealed conservation of viral genes across different host species, countries, and sampling years, suggesting a conserved cellular receptor and wide-ranging occurrence of bat influenza A viruses.

Influenza A viruses are major causes of human disease and are predominantly maintained in avian reservoirs (1). The segmented influenza A genome facilitates reassortment events in birds or intermediate hosts, such as swine and horses, leading to emergence of new variants potentially capable of causing zoonotic infections (2). Bats are major sources of zoonotic pathogens (3). In pioneering studies from 2012 and 2013, the first bat influenza A viruses,

Author affiliations: Charité-Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Institute of Virology, Berlin, Germany (A.C.A. Campos, L.G.B. Góes, A. Moreira-Soto, A.-L. Sander, C. Fischer, J.F. Drexler); Universidade de São Paulo-USP, Instituto de Ciências Biomédicas-ICB, São Paulo, Brazil (A.C.A. Campos, L.G.B. Góes, E.L. Durigon); Universidade Estadual Paulista Faculdade de Medicina Veterinária de Araçatuba, Araçatuba, Brazil (C. de Carvalho, W.A. Pedro, L.H. Queiroz); Universidade Estadual Paulista, Instituto de Biociências, Rio Claro, Brazil (G. Ambar, A.P. Cruz-Neto); Centro de Controle de Zoonoses, São Paulo (A.R. da Rosa, D.C. de Oliveira, L.F.A. Martorelli, A.P.G. Kataoka); German Centre for Infection Research, Germany (J.F. Drexler); Martsinovskiy Institute of Medical Parasitology, Tropical and Vector-Borne Diseases, Sechenov University, Moscow, Russia (J.F. Drexler)

DOI: <https://doi.org/10.3201/eid2502.181246>

termed H17N10 and H18N11, were discovered in 2 bat species, *Sturnira lilium* (little yellow-shouldered bat) and *Artibeus planirostris* (flat-faced fruit-eating bat) (4,5).

Bat-associated influenza A viruses are phylogenetically highly divergent from avian-associated influenza A viruses in their hemagglutinin (HA) and neuraminidase (NA) genes, suggesting these viruses represent ancient influenza A strains (2). Consistent with their genetic divergence, bat-associated influenza A surface proteins lack typical hemagglutination and neuraminidase activities (6), leading to the terminology HA-like (HL) and neuraminidase-like (NL) for bat-associated influenza surface proteins.

So far, only 4 individual bat specimens yielded influenza A genomic sequences during the pivotal investigations (4,5). HL18NL11 has only been found in 1 *A. planirostris* bat captured in Peru in 2010 (5), challenging definite host assessments. To investigate bat influenza A virus epidemiology, we investigated bats in southern Brazil during 2010–2014.

The Study

For this study, we sampled 533 individual bats representing 26 species and 3 families across 28 sampling sites (Table 1). Bats were captured using mist nets, euthanized, and necropsied and were identified on the basis of morphological criteria by trained field biologists as described previously (7). Only intestine samples were available for virological analyses. The Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais (21748–1), Instituto Ambiental do Paraná (235/10), and the ethics committee of the Institute of Biomedical Science from the University of São Paulo (56–18–03/2014) authorized sampling.

We tested intestine specimens from all bats using 2 highly sensitive, broadly reactive nested reverse transcription PCRs targeting different regions of the influenza A polymerase basic (PB) 1 gene (5,8). Positive results on both tests came from only 2 samples, from *Artibeus lituratus* great fruit-eating bats captured on March 7 and March 12, 2012, at 2 locations separated by 12 km in an Atlantic rainforest patch. No other sample was positive, yielding a 10.0% (2/20) overall detection rate in this site and 16.7% (2/12) detection in *A. lituratus* bats from this site (Table 1; Figure 1, panel A). Neither bat testing positive for influenza A virus showed signs of disease.

¹These senior authors contributed equally to this article.

Table 1. Bat species screened for influenza A virus, Brazil, 2010–2014*

Species	Family	No. samples	No. (%) PCR positive	Sampling site	Sampling years
<i>Artibeus fimbriatus</i>	Phyllostomidae	3	0	Iguaçu	2012
<i>Artibeus lituratus</i>	Phyllostomidae	129	2 (1.6)	Iguaçu , Central Paraná state, São Paulo cities	2010, 2011, 2012 , 2013, 2014
<i>Artibeus obscurus</i>	Phyllostomidae	1	0	São Paulo cities	2013
<i>Artibeus planirostris</i>	Phyllostomidae	4	0	Iguaçu, Central Paraná state, São Paulo cities	2010, 2012, 2014
<i>Carollia perspicillata</i>	Phyllostomidae	44	0	Iguaçu, Central Paraná state	2010–2012
<i>Cynomops planirostris</i>	Molossidae	6	0	São Paulo cities	2014
<i>Desmodus rotundus</i>	Phyllostomidae	15	0	São Paulo cities	2014
<i>Eptesicus furinalis</i>	Vespertilionidae	8	0	São Paulo cities	2013–2015
<i>Eumops auripendulus</i>	Molossidae	1	0	São Paulo cities	2014
<i>Eumops glaucinus</i>	Molossidae	44	0	São Paulo cities	2013–2015
<i>Eumops perotis</i>	Molossidae	8	0	São Paulo cities	2014–2015
<i>Glossophaga soricina</i>	Phyllostomidae	27	0	São Paulo cities	2013–2015
<i>Lasiurus cinereus</i>	Vespertilionidae	1	0	São Paulo cities	2013
<i>Lasiurus ega</i>	Vespertilionidae	1	0	São Paulo cities	2014
<i>Molossus molossus</i>	Molossidae	115	0	São Paulo cities	2013–2015
<i>Molossus rufus</i>	Molossidae	63	0	São Paulo cities	2013–2015
<i>Myotis nigricans</i>	Vespertilionidae	13	0	São Paulo cities	2013–2015
<i>Myotis riparius</i>	Vespertilionidae	1	0	São Paulo cities	2013
<i>Nyctinomops laticaudatus</i>	Molossidae	3	0	São Paulo cities	2014–2015
<i>Nyctinomops macrotis</i>	Molossidae	1	0	São Paulo cities	2014
<i>Phyllostomus discolor</i>	Phyllostomidae	2	0	São Paulo cities	2014
<i>Platyrrhinus lineatus</i>	Phyllostomidae	4	0	São Paulo cities	2014
<i>Promops nasutus</i>	Molossidae	1	0	São Paulo cities	2014
<i>Sturnira lilium</i>	Phyllostomidae	28	0	Iguaçu, Central Paraná state	2010–2012
<i>Tadarida brasiliensis</i>	Molossidae	9	0	São Paulo cities	2014
<i>Vampyressa pusila</i>	Phyllostomidae	1	0	Central Paraná state	2012
Total		533	2 (0.4)		

*Sampling sites were Parque Nacional do Iguaçu, Atlantic rainforest in western Paraná (Iguaçu); 26 cities across São Paulo state (São Paulo cities); and forest fragment in Paraná state (Central Paraná state). Bold indicates the site and year in which bats were captured that tested positive for influenza A virus.

A. lituratus bats were the most abundantly sampled species (Table 1).


The low overall influenza virus detection rate in this study (0.4%, 95% CI 0.0%–1.5%) was not significantly different by Fisher exact test from the previous 2 studies (1/110 bats for HL18NL11 [0.9%, 95% CI 0.0%–5.5%; $p = 0.43$]; 3/316 bats for HL17NL10 [1.0%, 95% CI 0.0%–2.9%; $p = 0.37$]). Apparently low rates of acute influenza A virus infection in bats are not consistent with high seroprevalence of 72% in different bat species according to a preliminary investigation (5) and may hint at seasonal variation in bat influenza virus infections, comparable to other batborne RNA viruses (9).

Sanger sequencing of the screening PCR amplicons suggested close genetic relatedness of the strains circulating in Brazil with the HL18NL11 strain circulating in Peru. Virus concentrations in the positive intestine specimens as determined by strain-specific quantitative real-time reverse transcription RT-PCR (Appendix Table 1, <https://wwwnc.cdc.gov/EID/article/25/2/18-1246-App.pdf>) were high (1.5×10^9 and 4.9×10^{10} RNA copies/g of tissue). High HL18NL11 concentrations in intestinal specimens are consistent with qualitative data from the pioneering study on HL18NL11 (5) and may suggest intestinal tropism and potential fecal shedding into the environment.

We determined the full coding sequence of all 8 segments of the viral genomes using primers aiming at amplifying overlapping regions of bat influenza A virus genomes (GenBank accession nos. MH682200–15) (Appendix Table 1). The 2 HL18NL11 variants in Brazil differed by 15 nt from each other across the combined 8 genomic segments. Four of those substitutions were nonsynonymous, causing amino acid exchanges in the PB2 (V203I), PB1 (R334K), nucleoprotein (G484S), and NA (V191I) genes (Table 2; Figure 1, panel B). This finding suggests recent common ancestry of the HL18NL11 variants identified in the 2 positive bats and was consistent with their detection in the same site 5 days apart. Comparison of the full coding sequence of the novel HL18NL11 variants revealed high sequence identity between the Peru and the Brazil strains, 93.5%–96.9% nucleotide identity across all 8 genomic segments (Table 2). The genomic relatedness of Peru and Brazil HL18NL11 strains was surprising given a time span of 2 years, a geographic distance exceeding 2,000 km, and 2 different bat species that tested positive in our study and the previous study (5).

All critical amino acid residues associated with influenza A virus replication and entry (4,5) were conserved between the Brazil and the Peru HL18NL11 strains, including the HA monobasic cleavage site motif

Figure 1. Bat influenza A(HL18NL11) virus detection and genomic characterization, Brazil, 2010–2014. A) Distribution of *Artibeus* species bats carrying HL18NL11 in Central and South America, according to the Red List of Threatened Species from the International Union for Conservation of Nature (https://www.iucnredlist.org). Orange star indicates the sampling site of an HL18NL11-positive bat in Peru (5); blue star indicates the sampling site of the HL18NL11-positive bats in Brazil for this study. Maps were created using QGIS2.14.3 (http://www.qgis.org) with data freely available from http://www.natureearthdata.com. B) Top, schematic representation of the genome organization of A/great fruit-eating bat/



Brazil/2301/2012 (HL18NL11) and amino acid exchanges (black lines) compared with A/great fruit-eating bat/Brazil/2344/2012 (HL18NL11) and Peru HL18NL11 (GenBank accession nos. CY125942–49). Nucleotide sequence identities between the concatenated HL18NL11 (Brazil), HL17NL10, and HL18NL11 (Guatemala and Peru) sequences were calculated in SSE version 1.2 (http://www.virus-evolution.org/Downloads/Software) with a sliding window of 200 and step size of 100 nt. C) Homology model of the HL protein of A/great fruit-eating bat/Brazil/2301/2012 viewed from the top, modeled on the published crystal structure retrieved from the SWISS-MODEL repository (https://www.swissmodel.expasy.org). The putative RBS is shown in blue, 3 highly conserved residues (W153, H183, and Y195) in HAs and HLs are in purple, and amino acid substitutions between Brazil strains and the Peru prototype strain are in red. D) Homology model of the NL of A/great fruit-eating bat/Brazil/2301/2012 viewed from the top, constructed as in panel C. The putative active site is shown in a blue circle, the 6 residues (R118, W178, S179, R224, E276 and E425) conserved in influenza A virus neuraminidase genes are in purple, and amino acid substitutions between Brazil strains and the Peru prototype strain are in red. HA, hemagglutinin; HL, hemagglutinin-like; NL, neuraminidase-like; RBS, receptor-binding site.

PIKETR/GLF (5). Thermodynamic modeling revealed that the amino acid exchanges observed between the Brazil and Peru HL18NL11 strains did not alter the three-dimensional structure of the HL and NL proteins, and neither mapped to the putative receptor binding site of the HL protein (Figure 1, panel C), nor to the putative active site of the NL protein (Figure 1, panel D) (6). This result suggests preservation of the biologic activity

of these glycoproteins in different bat species and supported a broadly conserved cellular receptor of bat influenza A viruses that differs from sialic acid receptors used by avian-associated influenza A viruses (10). Significantly fewer amino acid exchanges were observed between the HL proteins of Brazil and Peru bat influenza virus than between the respective NL proteins ($p = 0.007$ by Fisher exact test) (Table 2). The apparently

Table 2. Comparison of influenza A(HL18NL11) strain found in bats in Brazil with prototype strains from Peru

Gene	Nucleotide sequence identity	Amino acid exchange site	
		A/great fruit-eating bat/Brazil/2301/2012 (HL18NL11a)	A/great fruit-eating bat/Brazil/2344/2012 (HL18NL11b)
PB2	93.6%	V76I, R471K, T473N, V478I, I559V, R574K, S631N	V76I, V203I , R471K, T473N, V478I, I559V, R574K, S631N
PB1	93.7%	V54I, T56V	V54I, T56V, R334K
PA	94.4%	T70A, R116K, D158N, V231I, T254S, I552V, R711G	T70A, R116K, D158N, V231I, T254S, I552V, R711G
HL	96.0%	N167T, F251L	N167T, F251L
NP	96.8%–96.9%	N20T, K350R, L357M, I380L, I387V	N20T, K350R, L357M, I380L, I387V, G484S
NL	93.5%	I11V, I15L, V82I, V200I, L254I, A264T, V284I, D332E, V378I, G382E	I11V, I15L, V82I, V191I , V200I, L254I, A264T, V284I, D332E, V378I, G382E
M	95.4%	None	None
NS1	94.4%	R57K	R57K

*Bold indicates amino acid exchanges occurring in only 1 of the 2 Brazil strains compared to the Peru prototype strain. HA, hemagglutinin; HL, HA-like; M, matrix; NA, neuraminidase; NL, neuraminidase-like; NS, nonstructural protein; PA, polymerase acidic; PB, polymerase basic.




Figure 2. Phylogenetic relationships between bat influenza A viruses from Brazil and reference viruses. Phylogenetic trees show comparison of the 8 segments of representative influenza A virus genomes (PB2, PB1, PA, HA/HL, NP, NA/ NL, M, NS) with A/great fruit-eating bat/Brazil/2301/2012 (HL18NL11a; GenBank accession nos. MH682200–7) and A/great fruit-eating bat/Brazil/2344/2012 (HL18NL11b; GenBank accession nos. MH682208–15), shown in red. Maximum-likelihood trees were inferred using a general time-reversible substitution model with a gamma distribution and invariant sites. Black dots represent bootstrap values $\geq 75\%$ (1,000 replicants). Trees were generally rooted using influenza B/Lee/1940 (GenBank accession nos. DQ792894–901) (data not shown). Trees were constructed by using MEGA 6.0 (<http://www.megasoftware.net>). HA, hemagglutinin; M, matrix; NA, neuraminidase; NS1, nonstructural protein 1; NP, nucleoprotein; PA, polymerase acidic; PB, polymerase basic. Scale bars indicate nucleotide substitutions per site.

low rate of nonsynonymous substitutions in the HL-encoding genes of bat influenza A virus variants was reminiscent of strong purifying selection acting on the hemagglutinin genes in avian-specific influenza A virus strains (11). This finding may suggest comparable evolutionary dynamics between chiropteran and avian reservoirs. Definite assessments will require considerably larger datasets of bat influenza A virus strains.

A. lituratus bats and *A. planirostris* bats, in which HL18NL11 was originally detected in Peru, represent closely related, yet genetically and morphologically clearly distinct bat species (12). The distribution of these bat species overlaps (Figure 1, panel A), potentially facilitating virus exchange across the populations. Phylogenetic analyses confirmed the close genetic relationship between Peru and Brazil HL18NL11 variants across all 8 segments (Figure 2; Appendix Table 2), suggesting lack of reassortment events according to the available data. Our data thus suggest host associations of HL18NL11 beyond the species level, comparable to genus-level host associations of other batborne RNA viruses such as coronaviruses (13).

Conclusions

The zoonotic potential of HL18NL11 is unclear, yet human-derived cell lines were susceptible to infection by chimeric vesicular stomatitis virus pseudotyped with HL18 (14). The abundance of *A. lituratus* bats within Latin America (Figure 1, panel A) may thus facilitate spillover infections into other vertebrates across an underrecognized geographic and host range. Finally, *Artibeus* spp. bats have been used previously for infection studies including viruses with evolutionary origins in bats, such as Middle East respiratory syndrome coronavirus (15). The relatively large body size of *A. lituratus* bats (≈ 65 g) and ease of keeping these bats under laboratory conditions may thus facilitate experimental infection studies for HL18NL11 to elucidate the exact sites of HL18NL11 replication, receptor usage, and mode of transmission.

Acknowledgments

We thank Mariana Cristine Pereira de Souza, Cairo Monteiro de Oliveira, and Luciano Matsumiya Thomazelli for laboratory support.

This work was supported by FAPESP (São Paulo Research Foundation) through grants 2017/20744-5, 2014/15090-8, 2014/16320-7, 2013/11006-0, and 2008/57687-0 and the European Union's Horizon 2020 research and innovation programme through the ZIKAlliance project (grant agreement no. 734548) to J.F.D.

About the Author


Dr. Campos is a postdoctorate researcher affiliated with the University of Sao Paulo and Charité-Universitätsmedizin Berlin. Her research focuses on emerging viruses from bats.

References

- Olsen B, Munster VJ, Wallensten A, Waldenström J, Osterhaus ADME, Fouchier RAM. Global patterns of influenza A virus in wild birds. *Science*. 2006;312:384–8. <http://dx.doi.org/10.1126/science.1122438>
- Brunotte L, Beer M, Horie M, Schwemmle M. Chiropteran influenza viruses: flu from bats or a relic from the past? *Curr Opin Virol*. 2016;16:114–9. <http://dx.doi.org/10.1016/j.coviro.2016.02.003>
- Olival KJ, Hosseini PR, Zambrana-Torrel C, Ross N, Bogich TL, Daszak P. Host and viral traits predict zoonotic spillover from mammals. *Nature*. 2017;546:646–50. <http://dx.doi.org/10.1038/nature22975>
- Tong S, Li Y, Rivaviller P, Conrardy C, Castillo DA, Chen LM, et al. A distinct lineage of influenza A virus from bats. *Proc Natl Acad Sci U S A*. 2012;109:4269–74. <http://dx.doi.org/10.1073/pnas.1116200109>
- Tong S, Zhu X, Li Y, Shi M, Zhang J, Bourgeois M, et al. New World bats harbor diverse influenza A viruses. *PLoS Pathog*. 2013;9:e1003657. <http://dx.doi.org/10.1371/journal.ppat.1003657>
- García-Sastre A. The neuraminidase of bat influenza viruses is not a neuraminidase. *Proc Natl Acad Sci U S A*. 2012;109:18635–6. <http://dx.doi.org/10.1073/pnas.1215857109>
- Goes LGB, Campos ACA, Carvalho C, et al. Genetic diversity of bats coronaviruses in the Atlantic Forest hotspot biome, Brazil. *Infect Genet Evol*. 2016;44:510–3.
- Anthony SJ, Islam A, Johnson C, Navarrete-Macias I, Liang E, Jain K, et al. Non-random patterns in viral diversity. *Nat Commun*. 2015;6:8147. <http://dx.doi.org/10.1038/ncomms9147>
- Drexler JF, Corman VM, Wegner T, Tateno AF, Zerbinati RM, Gloza-Rausch F, et al. Amplification of emerging viruses in a bat colony. *Emerg Infect Dis*. 2011;17:449–56. <http://dx.doi.org/10.3201/eid1703.100526>
- Zhu X, Yu W, McBride R, Li Y, Chen LM, Donis RO, et al. Hemagglutinin homologue from H17N10 bat influenza virus exhibits divergent receptor-binding and pH-dependent fusion activities. *Proc Natl Acad Sci U S A*. 2013;110:1458–63. <http://dx.doi.org/10.1073/pnas.1218509110>
- Rejmanek D, Hosseini PR, Mazet JAK, Daszak P, Goldstein T. Evolutionary dynamics and global diversity of influenza A virus. *J Virol*. 2015;89:10993–1001. <http://dx.doi.org/10.1128/JVI.01573-15>
- Larsen PA, Marchán-Rivadeneira MR, Baker RW. Speciation dynamics of the fruit-eating bats (genus *Artibeus*): with evidence of ecological divergence in Central American populations. In: Adams RA, Pedersen SC, editors. *Bat evolution, ecology, and conservation*. New York: Springer Science + Business Media; 2013. p. 315–339. <https://doi.org/10.1007/978-1-4614-7397-8>
- Drexler JF, Corman VM, Drosten C. Ecology, evolution, and classification of bat coronaviruses in the aftermath of SARS. *Antiviral Res*. 2014;101:45–56. <http://dx.doi.org/10.1016/j.antiviral.2013.10.013>
- Moreira EA, Locher S, Kolesnikova L, Bolte H, Aydilto T, García-Sastre A, et al. Synthetically derived bat influenza A-like viruses reveal a cell type- but not species-specific tropism. *Proc Natl Acad Sci U S A*. 2016;113:12797–802. <http://dx.doi.org/10.1073/pnas.1608821113>
- Munster VJ, Adney DR, van Doremalen N, Brown VR, Miazgowiec KL, Milne-Price S, et al. Replication and shedding of MERS-CoV in Jamaican fruit bats (*Artibeus jamaicensis*). *Sci Rep*. 2016;6:21878. <http://dx.doi.org/10.1038/srep21878>

Address for correspondence: Jan Felix Drexler, Helmut-Ruska-Haus, Institute of Virology, Campus Charité Mitte, Charitéplatz 1, 10098 Berlin, Germany; email: felix.drexler@charite.de

EID Podcast: Bat Flight and Zoonotic Viruses



Bats are sources of high viral diversity and high-profile zoonotic viruses worldwide. Although apparently not pathogenic in their reservoir hosts, some viruses from bats severely affect other mammals, including humans. Examples include severe acute respiratory syndrome coronaviruses, Ebola and Marburg viruses, and Nipah and Hendra viruses.

Factors underlying high viral diversity in bats are the subject of speculation. The hypothesis is that flight, a factor common to all bats but to no other mammals, provides an intensive selective force for coexistence with viral parasites through a daily cycle that elevates metabolism and body temperature analogous to the febrile response in other mammals.

Visit our website to listen:

<http://www2c.cdc.gov/podcasts/player.asp?f=8632573>

**EMERGING
INFECTIOUS DISEASES**

Bat Influenza A(HL18NL11) Virus in Fruit Bats, Brazil

Appendix

RNA was extracted from 30 mg of tissue using the RNeasy Kit (QIAGEN, www.qiagen.com), followed by random hexamer-driven cDNA generation using the Superscript III reverse transcription kit (Thermo Scientific, www.thermofisher.com). Reactions were set up in a final volume of 20 μ L with 10 μ L of total RNA, 0.6 μ M of primers, 1x First-Strand Buffer, 0.5 mM (each) dNTP, 3.3 mM DTT, 1 μ g BSA, 40 U of Rnase OUT and 200 U SuperScript III. Hemi-Nested PCRs were performed in 25 μ L reactions with 1 μ L of cDNA (for first rounds) or PCR template (for second rounds), 1 μ M of each primer, 2.0 mM $MgCl_2$, 0.2 mM (each) dNTP and 1 U Platinum Taq Polymerase (Thermo Scientific). Thermocycling included a touchdown protocol with 94°C/3 min, 94°C/15 s, 68°C/30 s ($-1^\circ C$ per cycle) and 72°C/1 min during the first 10 cycles, followed by 45 cycles of 94°C/15 s, 58°C/30 s, 72°C/90 s, and 72°C for 7 min. One-step real-time RT-PCR-based quantification was done using primers targeting the PB1 gene and performed in 25 μ L reaction volumes with 5 μ L of RNA, 2.0 mM $MgCl_2$, 0.2 mM (each) dNTP, 0.4 μ M of each primer, 0.3 μ M of probe, and 1x PCR buffer with OneStep SSIII/Taq Enzyme Mix (Thermo Scientific). Amplification involved 55°C for 20 min (RT), followed by 94°C/3 min and 45 cycles of 94°C/15 s and 58°C/30 s with fluorescence read at the 58°C step, cooling at 40°C for 30 s on a LightCycler 480 thermocycler (Roche, www.roche.com). Quantification relied on photometrically quantified cRNA transcribed using the Megascript kit (Asuragen, www.asuragen.com) from a pCR4 vector containing the PCR target region (Thermo Scientific).

Appendix Table 1. Primers used for genomic amplification and sequencing

Gene	Primer name and position	Sequence 5' - 3'
PB2	BatFluPB2F1	AGCAGAAGCAGGTCARAGATTG
	BatFluPB2F2-632	TGGTTGCATACATGCTKGAAAGG
	BatFluPB2F3-1333	AGRCATTTCCAAAARGACTC
	BatFluPB2F4-1542	AAATGAAAAGGGAGAAYTWCT
	BatFluPB2F5-1336	CATTTCCAAAAGACTCTGGAG
	BatFluPB2F6-677	GGTTCCTGCCAGTTGCAGG
	BatFluPB2F7-727	CACCTAACCCAAGGCACGTG
	BatFluPB2F8-2089	GAGTCAGCAGTGTTAAGAGG
	BatFluPB2F9-2160	CGCAGAGCTGGATAAATTAGG
	BatFluPB2R7-763	ATTCTCWGCTTCMCCTCCTGG
	BatFluPB2R8-746	CCTGGWGTRTAYTGTGCTCCC

Gene	Primer name and position	Sequence 5'- 3'
	BatFluPB2R9-1678	TCCCAGTTYTTTAGTATCCAGTG
	BatFluPB2R11-1483	ATTCTYTCGTTGAAAGAGTATTCATC
	BatFluPB2R10-1762	CCYTTGGGKATTAAGTTTTGAAAWGG
	BatFluPB2R12-2256	GCTGTCTGGCTATCAGTAAGT
	BatFluPB2R13-1426	CCGGTAGTATCCCAATAATTCC
	BatFluPB2R14-1474	GACTTATTCTTATCCCAACCAAGTG
	BatFluPB2R1-2314	AGTAGAAAACAAGGTCATTTTTAGTG
	BRbatFluPB2F10-1000	GCTTTGGAGGCTATAACTTTAAG
	BRbatFluPB2F11-566	GATGCACAATTAGCGATCACCC
	BRbatFluPB2F12-627	CCAATTATGGTTGCATACATGC
	BRbatFluPB2R15-1572	TGCTTCACTTACTTCTTCTGG
	BRbatFluPB2R16-1611	GGGATGAATTGTAGTTTATTGG
	BRbatFluPB2R17-1066	GTTAATGTTTGAAGGTTTCCAG
PB1	BatFluPB1F1	AGCAGAAGCAGGCAAACACTATT
	BatFluPB1F2-1242	AGGRATGATGATGGGVATGTTT
	BatFluPB1F3-711	AARGAYGCAGAGAGAGGGWAAA
	BatFluPB1F4-1593	ATGAGYATAGGMACAACAGT
	BatFluPB1F5-1923	CACCATATGGARGTIGAAAGCAC
	BatFluPB1F6-1625	CATGATMAACAATGATCTAG
	BatFluPB1F7-1251	ATGGGSATGTTCAATATGC
	BatFluPB1F8-1594	TGAGTATAGGMACAACAG
	BatFluPB1F12-1926	CCATATGGAAGTCGAAAGCAC
	BatFluPB1F13-1958	TAATAATGCCAGCCCACG
	BatFluPB1F14-1961	TAATGCCAGCCCACGGGCCA
	BatFluPB1R3-759	CAAYCCTCTDATTTCATTCC
	BatFluPB1R4-1007	TGAACCATTCMGTTGICCT
	BatFluPB1R5-2012	GGTGTCCATGAATGAGTTG
	BatFluPB1R6-1007	TGAACCATTCMGTTGICCT
	BatFluR1PB1- 2322M2M1-1010	AGTAGAAAACAAGGCATTT
	BRbatFluPB1F10-1574	GAAACAATGAGTCTGCTGA
	BRbatFluPB1F11-1594	ATGAGTATAGGCACAACAG
	BRbatFluPB1R5-1424	CTACCAATTTGCAGATTCATAG
	BRbatFluPB1R7-1076	CATGTATCCCCGTCCAAGTC
PA	BatFluPAF1	AGCAGAAGCAGGTACTTARAC
	BatFluPAF2-680	CCWCCATCATTCAAGGACTAT
	BatFluPAF3-1144	GGAYTTTGAAGATTGTAAAG
	BatFluPAF4-1266	CTCAAACCTGGATYGAATTTGATG
	BatFluPAF6-1916	TGCAGAGTTCTTCTAGC
	BatFluPAF7-1943	TTCAACAGCATATATGCT
	BatFluPAF8	GCAGGTACTTARACAATGGAGAA
	BatFluPAR7-1176	GGWTCACTTTTGTATTGGAACA
	BatFluPAR8-1738	CATCTTCTCATTTCCATTCCC
	BatFluPAR9-1457	GTCTCTACATTTTRATTATTGG
	BatFluPAR10-1438	TGGKATYACYTGATATTCCTCCA
	BatFluPAR11-2089	GACCCAAGGATCATTAAATG
	BatFluPAR12-273	CAATAGTCCAGGCAACATTTT
	BatFluPAR13-300	CTATATTGGTCATGTTGCATATTG
	BatFluPAR1-2198	AGTAGAAAACAAGGTACTTT
HL	FluHaF1	AGCAGAAGCAGGGTSAYTATTAYTC
	BatFluHaF2-946	ACAGYACMCTGCCYTTTCA
	BatFluHaF3-991	AYTGTCCYAAATATGTGAARGC
	BatFluHaF4-756	GAGTTGTCAATCCTAATCAGAATC
	BatFluHaF5-951	CATCAAAATGCGATTGGAGATTGC
	BatFluHaF6-1247	AACTGCCAAAGAATCAAC
	BatFluHaR4-1305	CCATCATCAACTCTGTCACTGAG
	BatFluHaR5-1104	CATCAATCAWYCCTTGCCATCC
	BatFluHaR6-1729	GATTGACATTAGCTAACAC
	BatFluHaR7-242	GTTTCCCATAAGCCATGCAGG
	BatFluHaR1-1775	AGTAGAAAACAAGGGTSTTT
NP	BatFluNPF1	AGCAGAAGCAGGGTAAATAATC
	BatFluNPF2-24	CACATTGTGACATTTAAAGATG
	BatFluNPF3-739	CCACAGAAAGCAATGGTTGA
	BatFluNPF4-511	AATGGACCCAAGRATGTGCTC
	BatFluNPF5-1349	ACTGACATGAGAAGTGA
	BatFluNPF6-1386	ATTCTGATCCCAAAGAC
	BatFluNPF7-1344	GAATCACTGACATGAGAAGTGA
	BatFluNPF8-1303	GAAAACAATAATGGCTGCA

Gene	Primer name and position	Sequence 5'- 3'
	BatFluR1NP-1541NEP-878	AGTAGAAACAAGGGTATT
	BatFluNPR5-1189	GTCTTGATTGCCCAATAATG
	BatFluNPR6-1482	GAAATAAGAACCCTCGTCATTC
	BatFluNPR7-251	GGTATTTGTTCCTTCTTTTCGTC
NL	BatFluNaF1	AGCAGAAGCAGGAGTTTTTMA
	BatFluNaF2-892	CAAATCTYTGGAAATGATGCCAA
	BatFluNaF3-900	TGGAATGATGCCAARAGRCC
	BatFluNaHL18F-623	TGTGAGCATCCTTTATGGAG
	BatFluNaHL17F-629	AACAGACACTTTCTCGGCCAGCA
	BatFluNaF8-1232	TTATGAATCACGTGATTG
	BatFluNaR8-310	ATGCTGCACAGATTCTCTC
	BRbatFluNaR8-231	TGTTGAAGCTGAGAAGCT
	BRbatFluNaR9-362	GTCAGCATTTGTTTCATCA
	BRbatFluNaR10-993	ATTTGTAGTGCAATTCC
	BatFluNaR12-1396	CAAGGAGTTTTTTCTTATACATC
	BatFluNaR13-1395	CAAGGAGTTTTTTCTTATACATCC
	NewBatFluNaF1	AGCAGAAGCAGGAGT
	newBatFluNaF5-497	CAGTGTAAGTTAGGAGACC
	newBatFluNaF6-932	CCAGTCCTTCACTTACAC
	newBatFluNaF7-949	CTTTCAAGGAGCCATGCTTG
	newBatFluNaR4-511	CAGGTGTTGGAGGGTCTC
	newBatFluNaR5-573	CACTGAAAGCCATCATG
	newBatFluNaR6-1101	TCCTTTCTTGGATCCTGG
	newBatFluNaR7-1044	ATCATGAAACCTTGGATTCC
	inselnBatFluNaF1-177	AGCTGTCCAACGGGACTTCTG
	inselnBatFluNaF2-820	GGAACATYTSCHGGCTGGAAG
	inselnBatFluNaF3-916	ACAAATCTYTGGAAATGATGCCAA
	inselnBatFluNaR1-1129	CAAATCCWTTCTKGGATCCTGG
	inselnBatFluNaR2-655	CWGTTATWATTYCTCCATAWAGGAT
	inselnBatFluNaR3-596	GACAGTCCACTGAAAGCCATC
	BRbatFluNaF9-541	TTGAAGCTGTTGGCTGGA
	BRbatFluNaF10-598	TGTCCGTTGCAGGAGACG
	BRbatFluNaF13-1020	AAGACAACAACAGAGGGAGA
	BRbatFluNaR8-231	TGTTGAAGCTGAGAAGCT
	BRbatFluNaR9-362	GTCAGCATTTGTTTCATCA
	BRbatFluNaR10-993	ATTTGTAGTGCAATTCC
	BatFluNaR12-1396	CAAGGAGTTTTTTCTTATACATC
	BatFluNaR13-1395	CAAGGAGTTTTTTCTTATACATCC
M2/M1	BatFluM2M1F1	AGCARAAGCAGGCATTATYCAA
	New BatFluM2M1F1	AGCARAAGCAGGCATTATYC
	BatFluM2M1F2-575	CACTGCHAARGCCATGGARCAA
	BatFluM2M1F3-621	GCTGAAGCAATGGAAATTGC
	BRbatFluM2F4-547	GACATGAAAACCGAATGGCAAC
	BRbatFluM2F5-631	TGGAATTGCTTCACAAG
	BatFluM2M1R2-741	ACCAGAARAGRATGGGAAT
	BatFluM2M1R3-687	CACCCAACAACCTCCAGTGG
	BatFluM2M1R4-752	CTGCATCTGGATTCCCATC
	BatFluM2M1R5-690	GGCCACTGGAAGTTGTTGG
NEP/NS1	BatFluNEPF1	AGCAGAAGCAGGGTATCTAAAG
	New BatFluNEPF1a	AGCAGAAGCAGGGTATCTAA
	BatFluNEPF2-16	TCTAAAGACATAATGGAAYC
	BatFluNEPF2-40	CCGACAACATATCGCATTTTCAG
	BatFluNEPF3-514	AACCCTCTGTCTTTTGTACAG
	BatFluNEPF4-529	GTTACAGGACATACTGGAGAG
	BatFluNEPR2-583	GGATTTGAATGGAATGATAAC
	BRbatFluNEPR3-241	CATAGTAAGGCATGGCATC
	BRbatFluNEPR4-327	GATCATAATCCAATTTCTG
PB1 quantitative real-time RT-PCR	FluBR-rtF1	TGCAGAAGAAGCTGAAAYACTATAAGCTT
	FluBR-rtR	TGAACATSCCCATCATCATTCC
	Probe FluBR-rtP	FAM-TYGATGGGACAGCRTCACTGAGCCC-BHQ1

*HL, hemagglutinin-like; M2/M1, matrix protein 2 and matrix protein 1; NL, neuraminidase-like; NP = nucleocapsid, NEP/NS1, nuclearexport protein and non-structural protein 1; PA, PB1, PB2, polymerase genes.

†Numbers in primer names indicate the first nucleotide targeted in the Peruvian HL18NL11 prototype strain. For the degenerated bases, R = G/A, Y = C/T, S = G/C, W = A/T, M = A/C, K = G/T, H = A/C/T, I = inosine. FAM, 6-carboxyfluorescein; BHQ1, Black Hole Quencher1.

Appendix Table 2. Representative viruses used in phylogenetic analysis of Brazilian bat influenza A (HL18NL11) virus

NA	NA	Collection date	PB2 gene	PB1 gene	PA gene	HA gene	NP gene	NA gene	M gene	NS gene
H1	N1	1978	CY020300	CY020299	CY020298	CY020293	CY020296	CY020295	CY020294	CY020297
H1	N6	1977 Aug 2	CY004465	CY004464	CY004463	CY004458	CY004461	CY004460	CY004459	CY004462
H2	N1	1990 Apr 18	CY005420	CY005419	CY005418	CY005413	CY005416	CY005415	CY005414	CY005417
H3	N8	1963	CY032300	CY032299	CY032298	CY032293	CY032296	CY032295	CY032294	CY032297
H3	N5	1999 Oct 7	CY060258	CY060259	CY060260	CY060261	CY060262	CY060263	CY060264	CY060265
H4	N4	1979	CY045270	CY045269	CY045268	CY045263	CY045266	CY045265	CY045264	CY045267
H5	N2	1984 Feb 9	CY005764	CY005763	CY005762	CY014640	CY005760	CY014641	CY005759	CY005761
H6	N1	1979 Jan 1	CY005671	CY005670	CY005669	CY014623	CY005667	CY014624	CY005666	CY005668
H6	N2	2004 Dec 5	CY045478	CY045477	CY045476	CY045471	CY045474	CY045473	CY045472	CY045475
H7	N1	1934	CY077417	CY077418	CY077419	CY077420	CY077421	CY077422	CY077423	CY077424
H7	N7	1977	CY036902	CY036901	CY036900	CY036895	CY036898	CY036897	CY036896	CY036899
H8	N4	1968	CY005831	CY014662	CY005830	CY014659	CY005829	CY014660	CY005828	CY014661
H9	N6	1988 May 17	CY004574	CY004573	CY004572	CY005934	CY004570	CY004569	CY004568	CY004571
H10	N8	1965	CY005800	CY005799	CY014645	CY014644	CY005797	CY005796	CY005795	CY005798
H11	N1	1986 Nov 6	CY017772	CY017771	CY017770	CY017765	CY017768	CY017767	CY017766	CY017769
H12	N1	1983 Aug 6	CY005350	CY005349	CY005348	CY006006	CY005346	CY005345	CY005344	CY005347
H13	N2	1986 Jun 1	CY003901	CY003900	CY003899	CY005914	CY003897	CY003896	CY003895	CY003898
H14	N5	1982	CY130101	CY130100	CY130099	CY130094	CY130097	CY130096	CY130095	CY130098
H15	N9	1983	CY005724	CY005723	CY005722	CY006033	CY005720	CY005719	CY005718	CY005721
H16	N3	1988 May 16	CY004567	CY004566	CY004565	CY005933	CY004563	CY014569	CY004562	CY004564
H17	N10	May 2009	CY103873	CY103874	CY103875	CY103876	CY103877	CY103878	CY103879	CY103880
H17	N10	May 2009	CY103881	CY103882	CY103883	CY103884	CY103885	CY103886	CY103887	CY103888
H17	N10	Sep 2010	CY103889	CY103890	CY103891	CY103892	CY103893	CY103894	CY103895	CY103896
H18	N11	2010	CY125942	CY125943	CY125944	CY125945	CY125946	CY125947	CY125948	CY125949
H18	N11	2012 Mar 7	MH682200	MH682201	MH682202	MH682203	MH682204	MH682205	MH682206	MH682207
H18	N11	2012 Mar 12	MH682208	MH682209	MH682210	MH682211	MH682212	MH682213	MH682214	MH682215

*The influenza B strain used as an outgroup was B/Lee/1940 (accession numbers DQ792894–901).