



30 High Burden Countries



Risk of infection with tuberculosis



10 million cases in 2018



1,2 million deaths

250 thousand among PLHIV

4/10 deaths in PLHIV are due to TB



EH!WOZA.com








bhp.org.bw
ehwoza.com

<https://www.who.int/tb/publications/global-report/en/>

*Adolescent
Advocacy
Can End TB*





Youth Declaration

Ensure that no young person is left behind


Recognize young people as agents of change

Vision

increase knowledge about TB



Create space for dialogue



Gain Perspective



Increase interest in science/research



Raise TB Champions



Key messages about TB and HIV

People infected with TB should ensure they stay in well-ventilated rooms and take medication on time

Don't spread the news but spread the help because discriminating infected people doesn't make their load any lighter

People infected should go for counseling and accept themselves the way they are

If you have TB do not isolate yourself from others

No discrimination

Prevention of TB: Open the windows, cover your mouth while coughing, wash your hands

Signs of TB: coughing up blood, sweating, loss of weight, chest pains, fever

Accept others for who they are

Parents should disclose to their children the reason why they are taking medication

It's everyone's responsibility to know their status voluntarily not by circumstances such as pregnancy.

We should not personalize the problem so that we find solutions

Always go for check-ups and take treatment as prescribed

We should never let our religious beliefs get in the way of our treatment

There is still a lot of stigma even during the 21st century