


RESEARCH ARTICLE

Insecticide resistance is mediated by multiple mechanisms in recently introduced *Aedes aegypti* from Madeira Island (Portugal)

Gonçalo Seixas¹, Linda Grigoraki², David Weetman³, José Luís Vicente¹, Ana Clara Silva⁴, João Pinto¹, John Vontas^{2,5}, Carla Alexandra Sousa^{1*}

1 Global Health and Tropical Medicine, Instituto de Higiene e Medicina Tropical, Universidade Nova de Lisboa, Lisboa, Portugal, **2** Institute of Molecular Biology and Biotechnology, Foundation of Research and Technology, Heraklion, Greece, **3** Department of Vector Biology, Liverpool School of Tropical Medicine, Liverpool, United Kingdom, **4** Departamento de Planeamento, Saúde e Administração Geral do Instituto de Administração da Saúde e Assuntos Sociais, IP-RAM, Funchal, Madeira, Portugal, **5** Department of Crop Science, Agricultural University of Athens, Athens, Greece

* casousa@ihmt.unl.pt



OPEN ACCESS

Citation: Seixas G, Grigoraki L, Weetman D, Vicente JL, Silva AC, Pinto J, et al. (2017) Insecticide resistance is mediated by multiple mechanisms in recently introduced *Aedes aegypti* from Madeira Island (Portugal). PLoS Negl Trop Dis 11(7): e0005799. <https://doi.org/10.1371/journal.pntd.0005799>

Editor: Audrey Lenhart, Centers for Disease Control and Prevention, UNITED STATES

Received: April 11, 2017

Accepted: July 11, 2017

Published: July 24, 2017

Copyright: © 2017 Seixas et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This study was supported by the following projects: PTDC/SAU-EPI/115853/2009 (FEDER - COMPETE Program and Fundação para a Ciência e Tecnologia); GHTM-UID/Multi/04413/2013; and by the European Union's Horizon 2020 Research and Innovation Programme under ZIKAlliance Grant Agreement no. 734548. GS was

Abstract

Background

Aedes aegypti is a major mosquito vector of arboviruses, including dengue, chikungunya and Zika. In 2005, *Ae. aegypti* was identified for the first time in Madeira Island. Despite an initial insecticide-based vector control program, the species expanded throughout the Southern coast of the island, suggesting the presence of insecticide resistance. Here, we characterized the insecticide resistance status and the underlying mechanisms of two populations of *Ae. aegypti* from Madeira Island, Funchal and Paúl do Mar.

Methodology/Principal findings

WHO susceptibility bioassays indicated resistance to cyfluthrin, permethrin, fenitrothion and bendiocarb. Use of synergists significantly increased mortality rates, and biochemical assays indicated elevated activities of detoxification enzymes, suggesting the importance of metabolic resistance. Microarray-based transcriptome analysis detected significant upregulation in both populations of nine cytochrome P450 oxidase genes (including four known pyrethroid metabolizing enzymes), the organophosphate metabolizer *CCEae3a*, Glutathione-S-transferases, and multiple putative cuticle proteins. Genotyping of knockdown resistance loci linked to pyrethroid resistance revealed fixation of the 1534C mutation, and presence with moderate frequencies of the V1016I mutation in each population.

Conclusions/Significance

Significant resistance to three major insecticide classes (pyrethroid, carbamate and organophosphate) is present in *Ae. aegypti* from Madeira Island, and appears to be mediated by multiple mechanisms. Implementation of appropriate resistance management strategies including rotation of insecticides with alternative modes of action, and methods other than

funded by a FCT PhD grant (SFRH/BD/98873/2013). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

chemical-based vector control are strongly advised to delay or reverse the spread of resistance and achieve efficient control.

Author summary

Aedes aegypti is the major mosquito vector of dengue, chikungunya and Zika worldwide. After its introduction in Madeira, it took a few years for the first dengue outbreak to occur in the region. Control strategies rely mostly on the use of insecticides but their efficiency is often being hampered by the ability of mosquitoes to resist to the compounds used. In fact, previous vector control programs using insecticides failed to eradicate, or even, to limit the spread of *Ae. aegypti* in Funchal, and now, the mosquito is widely distributed throughout the southern coast of the island. Bioassays to determine insecticide susceptibility profiles were carried-out in two populations of Madeira Island and the molecular mechanisms underlying the observed insecticide resistance phenotype were investigated. Transcription levels of detoxification genes were analysed, and screenings for *kdr* mutations, V1016I and F1534C, associated with pyrethroid resistance were performed. Our study showed the up-regulation of several detoxification genes of multiple enzyme families associated with metabolic resistance, and the presence of the two *kdr* mutations, with the F1534C being fixed. Another suggested mechanism probably involved in the resistance phenotype is cuticle thickening, as several cuticle genes were found overexpressed. This study reinforces the importance of alternative control strategies to suppress *Ae. aegypti* population and thus reduce the likelihood of arbovirus transmission in the region.

Introduction

Aedes aegypti (Linnaeus, 1762) is the most important vector of dengue, chikungunya and Zika viruses. Originally from Africa, this mosquito species invaded other territories and currently has a worldwide tropical distribution, probably aided by globalization [1]. Of major concern is that novel arboviral outbreaks have occurred following the establishment of *Ae. aegypti* in new areas [2,3]. In Europe, the most recent example of invasion and subsequent arboviral outbreak is the case of Madeira Island, a Portuguese territory in the Atlantic Ocean. In 2005, *Ae. aegypti* was first recorded in the Santa Luzia parish of Funchal (the capital city of Madeira) and soon the vector spread into other parishes of Funchal [4,5]. Control measures based on source reduction were soon implement together with insecticide spraying mainly with pyrethroid insecticides, and application of *Bacillus thuringiensis israelensis* (*Bti*) for larval control [6,7]. This insecticide-based strategy was applied only in Santa Luzia municipality of Funchal, during the years 2006–2008. This strategy included outdoor ULV applications with alfa-cypermethrin and indoor sprayings with tetramethrin and d-fenothrin, in households, and pyrethrins, in warehouses and similar facilities. A liquid formulation of *Bti* was used to treat water fountains or other breeding sites that had standing water [7]. Despite the vector control measures, *Ae. aegypti* continued to expand throughout the entire south coast of the island [6].

In 2012, a dengue outbreak was declared by the Madeira Health Authorities with a total of 2,168 cases reported from September 2012 to March 2013 [7]. In this period, a total of 78 imported dengue cases were notified in 13 European countries, consisting of travelers that had visited the island during the outbreak [7]. This event was of particular public health concern given that competent dengue vector *Aedes albopictus* populations are established in southern

Europe and have been implicated in autochthonous transmission of dengue and chikungunya viruses in Italy, France and Croatia [8–10].

Insecticide-based vector control efforts did not succeed in reducing the mosquito population, which thrived along the southern coast of the island. This is the most densely inhabited part of the island, resulting in a great availability of human hosts and breeding sites, which coupled with favourable climatic conditions may explain the rapid establishment of *Ae. aegypti*. Insecticide resistance might also have contributed to the apparently limited effectiveness of the implemented insecticide-based control measures, but investigation to date has been limited.

Reduced susceptibility to insecticides has been reported frequently in *Ae. aegypti* and is primarily associated with two major mechanisms of insecticide resistance, metabolic detoxification and target-site mutations. Over-expression of genes belonging to esterase, Glutathione-S-transferase (GST) and cytochrome P450 oxidase detoxification enzyme families has been reported in insecticide resistant *Ae. aegypti* populations from a broad range of geographic locations (reviewed in [11,12]). The role of several detoxification enzymes in insecticide resistance, including six P450s in pyrethroid resistance and the esterase *CCEae3a* in organophosphate resistance has been confirmed by *in vitro* and/or *in vivo* functional validation studies [13,14].

Multiple point mutations have been identified in the voltage-gated sodium channel gene (*Vgsc*) of pyrethroid-resistant *Ae. aegypti* populations worldwide. Of these, mutations at two codons are most commonly involved in resistance to pyrethroids in *Ae. aegypti*, V1016G or I and F1534C, which may act multiplicatively, especially in combination with an additional mutation, S989P (reviewed in [12]).

Cuticle thickening has also been implicated in insecticide resistance by interfering with the penetration of the insecticide and thus with the amount and rate of insecticide that reaches its target-site [15]. This type of resistance has been documented in several insect species including disease vectors [16–18], and genes encoding cuticle proteins have been found to be over-expressed in insecticide resistance strains of *Ae. aegypti* and *Ae. albopictus*, as well as *Anopheles stephensi*, and *Culex pipiens pallens* [19–22].

A previous study [23] detected the V1016I and F1534C point mutations in the *Vgsc* gene of *Ae. aegypti* from Madeira suggesting that target-site resistance is present on the island. However, no information on the prevalence of insecticide susceptibility, nor on other mechanisms of resistance in Madeira are available to date.

In order to clarify possible causes of inefficacy of insecticide-based control measures implemented on the island, and to assist health authorities in the planning of new interventions, we characterised resistance phenotypes and underlying mechanisms in *Ae. aegypti* from Madeira.

Methods

Study site and mosquito sampling

The archipelago of Madeira comprises seven islands and two islets situated in the north Atlantic *ca.* 685 km off the coast of Morocco, West Africa. Madeira is the largest island with 742 km² and a population of around 270,000 inhabitants. The island has a mountainous topography and most of the population lives along the coast, especially in the south, where the capital Funchal is located (32°39'4"N 16°54'35"W) and where nearly half of the population lives. This area is densely populated and domestic flower pots, which are the major breeding sites for *Ae. aegypti* in the island, are very abundant [24]. The subtropical climate, hilly landscape with exuberant Laurissilva forest and relative proximity to the European continent make this insular territory a popular tourist destination [25].

Aedes aegypti were sampled by ovitrap collections carried out in Funchal and Paúl do Mar between September and November 2013. Ovitrap distribution in both localities is available in

the citizen science online platform (<http://iasaude.sras.gov-madeira.pt/naomosquito/>). No specific permits were required for the described field collections. Immatures were reared to adulthood for subsequent use in insecticide susceptibility bioassays. Mosquito rearing and bioassays were performed in the facilities of Direção Regional da Agricultura, provided through a cooperation agreement with the Instituto de Administração da Saúde e Assuntos Sociais (IASAUDE). The *Ae. aegypti* Rockefeller strain was used as susceptible reference colony. Mosquitoes were reared in a controlled environment with stable temperature ($26\pm 2^{\circ}\text{C}$), relative humidity ($70\pm 5\%$) and photoperiod (12h/12h light/dark). A subset of non-insecticide exposed females, which emerged from field collected immatures and from the Rockefeller reference susceptible strain were frozen in liquid nitrogen for subsequent use in biochemical assays. In addition, individuals from the Funchal and Paul do Mar populations, as well as individuals from the susceptible laboratory colonies Rockefeller and New Orleans were stored in RNALater (Invitrogen) to be used in the gene expression analysis.

Susceptibility bioassays

Bioassays were carried out with 3–5 day old non-blood fed females using WHO insecticide susceptibility tests and protocols [26,27]. Filter papers impregnated with insecticide (bendiocarb 0.1%, cyfluthrin 0.15%, fenitrothion 1.0% and permethrin 0.75%) were provided by WHO-University of Sains Malaysia (Penang, Malaysia). Insecticides were chosen according to the three main classes of chemicals allowed to be used in vector control in the region, and based on previous knowledge regarding insecticide resistance status of the *Ae. aegypti* populations [7]. Females were exposed to the insecticide in groups of 20–25 per tube, for one hour. Four or five replicates per insecticide were used, depending in mosquito availability. After exposure, mosquitoes were transferred to a holding tube and supplied with a 10% sugar solution on a cotton pad. Mortality was scored 24 hours after exposure. The susceptibility status of each mosquito population was assessed according to WHO recommendations, in which a mosquito population is deemed resistant to a given insecticide if mortality rates are below 90% [26,27] when testing a minimum of 100 specimens. The surviving mosquitoes (considered resistant) and dead (considered susceptible) were stored individually in 1.5ml tubes filled with silica gel desiccant for DNA-based analysis.

In addition, WHO susceptibility tests were carried out with pre-exposure to synergists to block the action of P450s, esterases and GSTs, in order to assess presence of metabolic resistance. Females were exposed to papers impregnated with 4% piperonyl butoxide (PBO) or 8% diethyl maleate (DEM) for one hour and then immediately exposed to each insecticide or to control papers as described above. Mortality rates were scored after 24 hours.

Biochemical assays

Biochemical assays were performed to quantify the enzymatic activity of the major detoxification families: esterases, Glutathione-S transferases (GST) and cytochrome P450 oxidases (MFO), following WHO protocols [28]. Forty 3–5 days- old individual females from Funchal, Paul do Mar and an equal number of the Rockefeller reference strain were used in each assay. Comparisons of enzyme activity between field and reference mosquitoes were tested using a Mann-Whitney non-parametric analysis using Graphpad Prism v 6.03.

Microarray: RNA extraction, labeling and hybridization

Gene expression analysis was carried out at the Liverpool School of Tropical Medicine, UK. Three day-old females, non-blood fed and not exposed to insecticides, that were F1 progeny of mosquitoes collected in Funchal and Paul do Mar were compared to three day-old non-blood

fed, insecticide unexposed females from the susceptible reference colony Rockefeller. In addition, the Funchal population was compared to the second susceptible reference colony New Orleans, to further reduce the possibility that differences observed in expression levels could be related to differences in the genetic background of the laboratory strains unrelated to phenotype. Total RNA was extracted from four replicate batches of five mosquitoes using the Arc-turus PicoPure RNA isolation kit (Applied Biosystems). In all cases, RNA was treated with DNase using the RNase-free DNase Set (Qiagen), according to the manufacturer's instructions. Quantity and quality of the RNA extracts were evaluated with a Nanodrop spectrophotometer (Nanodrop Technologies) and a 2100 Bioanalyzer (Agilent Technologies), respectively. The RNA pools were amplified and labeled using the Low Input Quick Amp Labeling Kit (Agilent Technologies). Quality and quantity of labeled cRNA was assessed as above before further use. Four hybridizations for each comparison (*i.e.* Funchal vs New Orleans, Funchal vs Rockefeller and Paul do Mar vs Rockefeller) were performed using the 15k Agilent "Aedes microarray" (ArrayExpress accession number A-MEXP-1966). After 17 hours of hybridization at 65°C, the array was washed to remove non-specifically bound probes, using Agilent microarray washing buffers. Scanning was performed immediately after washing on an Agilent G2205B microarray scanner.

Microarray data analysis

Data processing was performed using the Agilent Feature extraction software and analysis of normalized data used Genespring v13. A strict filtering criterion was used for inclusion of probes where all had to be detectable (or marginal) in every array across each dataset, resulting in data from 9083 acceptable probes. Probability of differential expression was determined by a one-sample *t*-test (null hypothesis of a ratio of field/colony sample expression of 1) with the *P*-value threshold set at $P < 0.05$. A fold change threshold of $FC > 2$, or $FC < -2$ (for underexpressed probes) was also implemented. We employed a replication criterion for significance, such that a gene was considered differentially expressed if the probability and fold-change thresholds were met for each of the three comparisons with the susceptible reference strains. Although individually the use of a threshold of $\alpha = 0.05$ would lead to a high expected number of false positives ($N \approx 450$, ignoring the additional FC criterion), the use of a strict 3/3 replication criterion reduces this dramatically to $N \approx 1$ [29], again ignoring additional stringency from the FC criterion. Owing to this strict replication procedure we also identified probes as potentially significant if they exhibited $P < 0.05$ in 2/3 analyses and an extreme level of expression ($FC > 20$). For representation but not assessment of significance, fold changes were averaged and *P*-values combined using Fisher's method for combining probabilities.

Microarray validation by qRT-PCR

The transcription level of candidate overexpressed genes was validated by qRT-PCR in the Funchal population. Two micrograms of DNase-treated RNA from each sample (four biological replicates for each strain: Funchal, New Orleans and Rockefeller) were reverse-transcribed using oligo(dT)₂₀ (Invitrogen) and Superscript III (Invitrogen). Amplification reactions of 25 µl final volume were performed in a MiniOpticon Two-Color Real-Time PCR Detection System (BioRad) using 2 µl of 1/25 diluted cDNA, 0.2 µM primers (S1 Table) and Kapa SYBR FAST qPCR Master Mix (Kapa-Biosystems). For normalization of results, the ribosomal proteins L8_AAEL000987 and S7_AAEL009496 were used [30]. A fivefold dilution series of pooled cDNA was used to assess the efficiency of the qPCR reaction for each gene specific primer pair. A no template control (NTC) was included to detect contamination, and a melting curve analysis was done to check for the presence of a unique PCR product. The thermal

profile of reactions was 95°C for 3min followed by 40 cycles of 95°C for 15sec, 58°C 30sec and 60°C for 30sec. Relative expression analysis was performed according to Pfaffl [31].

DNA isolation and *kdr* genotyping

A subsample of mosquitoes phenotyped as susceptible or resistant by WHO assays to pyrethroid insecticides (without pre-exposure to synergists) were genotyped for the presence of the two previously-detected mutations in the *Vgsc* gene [23]. Genomic DNA was extracted according to Collins et al [32]. Two allele-specific PCR assays (AS-PCR) were used to genotype *kdr* mutations V1016I and F1534C [23]. For the V1016I mutation, the protocol used was adapted from Saavedra-Rodriguez et al [33]. Amplifications were carried out in 25 µl of reaction mixture containing 1X buffer, 3 mM of MgCl₂, 0.2 mM of each dNTP, 0.1 µM of primers Val1016f, Iso1016f and Iso1016r and 1U of *Taq* DNA polymerase. The PCR conditions were identical to those described in Saavedra-Rodriguez et al [33]. PCR products were separated by electrophoresis (90 minutes at 90V) in an ethidium bromide-stained 3% agarose gel and photographed under UV light.

The tetra-primer PCR assay described in Harris et al [34] was used to genotype the F1534C mutation. Each reaction of 25 µl contained 1X PCR buffer, 2.5 mM MgCl₂, 0.4 mM of each dNTP, 0.25 µM of primers AaEx31P, AaEx31Q, AaEx31wt and AaEx31mut and 1 U of *Taq* DNA polymerase. The cycling conditions were the same used in Harris et al [34]. PCR products were size-fractionated by electrophoresis in ethidium bromide stained 2% agarose gels at 100V (45 minutes) and photographed under UV light.

All PCR assays contained negative controls (*i.e.* no DNA template) and positive controls, consisting of samples of known genotype confirmed by DNA sequencing [23].

Results

Susceptibility bioassays

Aedes aegypti from Funchal were found to be resistant to all insecticides tested (Fig 1A), with mortality rates ranging between 10.9% (after permethrin exposure) and 77.5% (after fenitrothion exposure). Mortality rates increased significantly when females were exposed to one or both of the synergists before the insecticide, suggesting involvement of metabolic resistance (Fig 1A). This was particularly evident for permethrin, after exposure to either PBO or DEM, and also for fenitrothion, for which complete restoration of susceptibility was attained with both synergists. For cyfluthrin and bendiocarb a significant increase in mortality was observed after exposure with PBO only.

Resistance to pyrethroids and carbamates was also observed in the Paúl do Mar population, with mortality rates between 2% (permethrin), 63.1% (bendiocarb) and 88% (cyfluthrin) (Fig 1B). In contrast, exposure to fenitrothion yielded 100% mortality indicating full susceptibility to this insecticide. As in the Funchal population, synergist assays suggest the presence of metabolic resistance (Fig 1B). Exposure to both synergists, before insecticide contact, led to a significant increase in mortality rates with permethrin. In addition, near-full susceptibility to bendiocarb was achieved when previously exposing these mosquitoes to PBO. Synergist assays with cyfluthrin, in this *Ae. aegypti* population, did significantly alter the mortality rate. No mortality was observed in the control mosquitoes whether exposed to control papers alone or to synergists (with no insecticide).

Biochemical assays

A significantly higher enzymatic activity was detected for both α - and β -esterases in both populations when compared to the susceptible Rockefeller reference strain (Mann-Whitney tests,




Fig 1. Susceptibility levels of *Ae. aegypti* from Funchal (A) and Paúl do Mar (B) to insecticides. * Significant differences in mortality rates between exposures with and without synergists (Fisher's exact test, $P < 0.05$). The error bars represent standard deviation.

<https://doi.org/10.1371/journal.pntd.0005799.g001>

$P < 0.05$), while no significant difference was observed in the enzymatic activity of GSTs. A statistically significant difference was seen in mixed function oxidases only in the Paúl do Mar population ($P = 0.01$). Funchal population did not show differences in the enzymatic activity of this enzyme family ($P = 0.7204$) (Fig 2).




Fig 2. Activity profile of esterases, GST and mixed function oxidases (MFO) enzyme families of *Ae. aegypti* from Funchal and Paúl do Mar. * Significant differences in enzymatic activities between the wild population and the Rockefeller susceptible strain (Mann-Whitney test, $P < 0.05$).

<https://doi.org/10.1371/journal.pntd.0005799.g002>

Microarray analysis of differential expression

From a total of 9083 probes passing quality control, 141 were differentially expressed ($|\text{Fold Change}| > 2$, $P < 0.05$, for three out of three comparisons to susceptible strains) across the Funchal and Paul do Mar populations (S1 Table). Among the 86 probes which were up-regulated (Fig 3), 11 genes were members of the three detoxification enzyme super-families (P450s, GSTs and Carboxyl/choline esterases) (Table 1). A further gene, *Cyp9J32* was also included in the table with the over-expressed genes as it showed extreme over-expression, and only marginally missed the threshold P-value in one (out of three) comparison. Thus we considered this as a false negative, resulting from our strict filtering procedure (Table 1). The P450 oxidases had the highest representation with nine genes, all from the CYP6 and CYP9 sub-families and included four known pyrethroid metabolizers, of which *Cyp9J32* and *Cyp9J28* were particularly strongly over-expressed ($\text{FC} > 20$). Other detoxification genes found overexpressed in both wild populations were the *GSTd4*, *GSTd1* and *CCEae3a*, which metabolizes temephos-oxon (the toxic form of the larvicide temephos) and thus may be of relevance for resistance in the populations, although we did not evaluate temephos resistance in this study.




Fig 3. Commonly up-regulated transcripts in *Ae. aegypti* populations from Funchal and Paúl do Mar. A rhombus shape is used for transcripts meeting the criteria Fold Change >2 and P<0.05 in all three comparisons performed. Among these genes are the rer1 protein (possibly involved in the retrieval of endoplasmic reticulum proteins), a lysosomal aspartic protease precursor and many genes with unknown function. A square shape is used for potentially significant up-regulated transcripts, exhibiting a P<0.05 in 2/3 analyses and an extreme level of expression (FC>20). All transcripts falling into the categories of detoxification genes, hexamerins or genes encoding for cuticular proteins are shown with red color. *Cyp9J27* is present in two distinct locations in the genome and so is represented twice, though with different accession numbers (Table 1).

<https://doi.org/10.1371/journal.pntd.0005799.g003>

Table 1. Commonly overexpressed transcripts in *Ae. aegypti* from Funchal and Paul do Mar belonging to detoxification gene families.

Class of detoxification gene	Gene accession number	Gene name	Funchal vs Rockefeller (FC)	P-value	Funchal vs New Orleans (FC)	P-value	Paul do Mar vs Rockefeller (FC)	P-value
P450s	AAEL008846	<i>Cyp9J32</i>	41	0.006	53	0.004	10	0.060
	AAEL014617	<i>Cyp9J28</i>	31	0.028	30	0.019	10	0.008
	AAEL014893	<i>Cyp6BB2</i>	15	0.005	19	7×10^{-4}	3.6	0.006
	AAEL014607	<i>Cyp9J27</i>	14	0.021	18	0.018	3.5	0.023
	AAEL014616	<i>Cyp9J27</i>	14	0.020	16	0.016	4.6	0.006
	AAEL001288	<i>Cyp9M5</i>	6.6	6×10^{-4}	8.5	0.009	2.3	0.001
	AAEL001312	<i>Cyp9M6</i>	5.4	0.015	4.7	0.006	4	0.031
	AAEL009124	<i>Cyp6N12</i>	5.1	3×10^{-4}	6.3	0.003	2.2	0.002
	AAEL017297	<i>Cyp6M9</i>	4.2	0.007	3.7	0.001	3	0.047
Esterases	AAEL005112	<i>CCEae3a</i>	2.7	0.009	3.4	0.012	2.1	0.001
GSTs	AAEL001054	<i>GSTd4</i>	24	4×10^{-4}	22	2×10^{-4}	9.8	0.004
	AAEL011741	<i>GSTs1</i>	4.9	2×10^{-4}	2.5	7×10^{-5}	2.2	0.032

FC represents the relative fold change in expression in the Funchal or Paul do Mar population compared to the respective susceptible colony. Bold type indicates a known insecticide metabolizer [13,14,40].

<https://doi.org/10.1371/journal.pntd.0005799.t001>

Beyond genes belonging to detoxification gene families the analysis revealed also the over-expression of eight transcripts encoding putative cuticle proteins, which have been implicated in resistance through lower insecticide penetration and also of three transcripts (AAEL011169, AAEL013759 and AAEL000765, which was highly and significantly, $P < 0.05$, up-regulated in 2/3 comparisons) encoding hexamerins, which are involved in cellular trafficking and have previously been linked to insecticide resistance [35] (Fig 3, S1 Table).

qRT-PCR validation

Quantitative real time PCR was used to validate the differential expression of five candidate genes detected as significantly up-regulated in the Funchal population (for which microarray data *vs.* both susceptible colonies was available) compared to the two susceptible colonies. We tested the highly overexpressed P450s *Cyp9J32* and *Cyp9J28*, the more highly overexpressed of the glutathione S transferases, *GSTd4*, the more highly overexpressed of the hexamerins, AAEL013757 and the more highly overexpressed member of genes encoding for putative cuticular proteins, AAEL002246 (S1 Table) (Table 2). Although the relative levels of overexpression estimated by qRT-PCR did not correspond closely with values obtained from the microarray experiment, the estimates from qRT-PCR confirmed up-regulation of the tested genes, thus providing validation of the significance indicated by microarrays.

Kdr genotyping

Genotyping of the *kdr* locus was performed on a total of 91 Funchal and 80 Paúl do Mar specimens that had been previously exposed to cyfluthrin or permethrin (Table 3). The 1534C mutation was found in every specimen genotyped suggesting it may be fixed in both populations, while the V1016I mutation showed moderate and similar frequencies ranging from 17% in Funchal to 23% in Paúl do Mar (Fisher's exact tests, $P = 0.263$). Owing to ubiquitous occurrence of the 1534C mutation testing association with resistance was not possible. Although the V1016I frequency was slightly higher in resistant mosquitoes when compared to susceptible ones, there was no significant association between *kdr* genotypes and the resistance phenotype for either insecticide (Fisher's exact tests, Funchal—cyfluthrin: $P = 0.491$; permethrin: $P = 0.699$; Paúl do Mar—cyfluthrin: $P = 0.316$; permethrin: $P = 0.219$).

Table 2. Validation of the transcriptional up-regulation of five candidate genes through qRT-PCR.

Transcript	Reference strain	qRT-PCR	Microarray
		Fold change (95% CI)	Fold Change
Hexamerin (AAEL013757)	Rockefeller	63 (36–90)	57
	New Orleans	123 (54–191)	97
<i>Cyp9J32</i> (AAEL008846)	Rockefeller	17 (13–21)	41
	New Orleans	107 (81–132)	53
Cuticular (AAEL002246)	Rockefeller	42 (16–69)	36
	New Orleans	65 (25–105)	51
<i>Cyp9J28</i> (AAEL014617)	Rockefeller	13 (6–21)	31
	New Orleans	104 (63–144)	30
<i>GSTd4</i> (AAEL001054)	Rockefeller	169 (34–305)	24
	New Orleans	268 (148–387)	22

The relative expression ratio of five candidate genes in the Funchal population compared to two susceptible laboratory colonies (New Orleans and Rockefeller) is shown. Values are estimated from four biological replicates and 95% confidence intervals are shown. Estimated values from the microarray experiment are given for comparison.

<https://doi.org/10.1371/journal.pntd.0005799.t002>

Table 3. Summary of *kdr* genotyping data in Funchal and Paúl do Mar *Ae. aegypti* populations.

Locality	Insecticide	N	V1016I				F1534C			
			V/V	V/I	I/I	F.(I)	F/F	F/C	C/C	F.(C)
Funchal	Cyfluthrin resistant	32	20	10	2	0.22	0	0	32	1.00
	Cyfluthrin susceptible	19	15	4	0	0.11	0	0	19	1.00
	Permethrin resistant	32	21	11	0	0.17	0	0	32	1.00
	Permethrin susceptible	8	6	2	0	0.13	0	0	8	1.00
	Total	91	62	27	2	0.17	0	0	91	1.00
Paúl do Mar	Cyfluthrin resistant	12	6	5	1	0.29	0	0	12	1.00
	Cyfluthrin susceptible	36	21	15	0	0.21	0	0	36	1.00
	Permethrin resistant	30	18	11	1	0.22	0	0	30	1.00
	Permethrin susceptible	2	0	2	0	0.5	0	0	2	1.00
	Total	80	45	33	2	0.23	0	0	80	1.00

N: sample size. Values correspond to absolute numbers for each genotype. F.(I) and F.(C) are the relative frequencies of the mutant allele for each mutation analyzed.

<https://doi.org/10.1371/journal.pntd.0005799.t003>

Discussion

The results of this study showed that *Ae. aegypti* from Madeira Island is resistant to insecticides of different chemical classes: carbamates (bendiocarb), organophosphates (fenitrothion) and both type I (permethrin) and type II (cyfluthrin) pyrethroids. Diagnostic exposures to these insecticides yielded mortality rates below the thresholds recommended by WHO to consider a mosquito population resistant [26,27], excepting the case of fenitrothion in Paúl do Mar. Combined bioassays with synergists and analysis of detoxification enzyme activities indicated the presence of enzyme-mediated metabolic resistance, and/or cuticular resistance. Pre-exposure to PBO, which inhibits P450s, some esterases and may also enhance cuticular penetration by the insecticide [36] resulted in a significant mortality increase for all insecticides tested. Synergist pre-exposures suggest involvement of the three major detoxification enzyme families in the resistance phenotype of *Ae. aegypti* from Madeira island. Biochemical assays only partially agreed with the results obtained by the bioassays with synergists, as significantly elevated enzymatic levels were detected for esterases only. These discrepancies were not completely unexpected as these assays cannot be considered reciprocal. While synergists act as inhibitors of enzymes suspected to be implicated in resistance, biochemical assays are a measure of enzyme activity without a direct link with the resistance phenotype. Furthermore, biochemical assays employ generic substrates which may not be recognized by all variants of these large enzyme families, resulting in reduced sensitivity and specificity [36, 37, 38]. The microarray-based transcriptomic analysis showed overexpression of genes belonging to the three major detoxification enzyme families in agreement with the bioassays with synergists. The majority of overexpressed detoxification genes were cytochrome P450 oxidases, including *Cyp9J32*, *Cyp9J28*, *Cyp9J27*, *Cyp6BB2* and *Cyp9M6*, which have been found overexpressed in pyrethroid resistant *Ae. aegypti* populations from multiple countries [12,36,39]. In particular, *Cyp9J28* is an efficient pyrethroid metabolizer [13] that has also been shown to confer reduced susceptibility to deltamethrin when ectopically expressed in *Drosophila melanogaster* [40]. *Cyp9J32*, the most prominent pyrethroid metabolizer in *Ae. aegypti*, showing a very high catalytic efficiency against pyrethroids [13], was the most overexpressed P450 gene. The carboxyl-esterase *CCEae3A*, which has previously been associated with resistance to the organophosphate temephos-oxon [14] in both *Ae. aegypti* [30] and *Ae. albopictus* [41], and acts via sequestration and metabolism of temephos [14], was also among the overexpressed genes in Funchal and Paúl do

Mar. This should be taken into account if temephos is considered for vector control in Madeira. The overexpression of two GSTs (*GSTd4* and *GSTs1*) was also revealed by the microarray analysis. Of these, *GSTd4* has been detected as highly overexpressed in the strongly permethrin resistant Singapore SP strain [42].

In addition to detoxification gene families, eight transcripts encoding putative cuticular proteins were up-regulated possibly indicating alteration of the cuticle as a mechanism of resistance. Resistance related to the cuticle by lowering the amount or rate of insecticide penetrating into the body has been reported in insects such as *Helicoverpa armigera* [43], *Drosophila melanogaster* [44], the *Trypanosoma cruzi* vector *Triatoma infestans* [16], and the *Plasmodium* vector *Anopheles funestus* [17]. Thickening of the whole cuticle, as well as the epicuticle layer, due to an increased number of cuticular hydrocarbons, has also been recently described in a multi-resistant strain of the malaria vector *Anopheles gambiae* [18]. Furthermore, there were also three transcripts (one marginally non-significant) encoding hexamerins among the most highly overexpressed genes. The link of hexamerins to insecticide resistance is poorly understood, but previous studies suggest a role for these storage proteins in cuticle formation [45,46]. It is also plausible that composition and thickening of the cuticle might reflect adaptive responses to environmental challenges and/or seasonality, rather than insecticide resistance. Seasonal cuticular variations, mostly associated with adaptation to aridity (desiccation tolerance), have been previously observed in other insect species, including scorpions [47], crickets [48] and more recently in the malaria vector *A. coluzzii* [49]. There is evidence that *Ae. aegypti* of Madeira derives from a tropical south American source population [23], thus, adaptation of this mosquito to the more temperate climate of Madeira might have involved changes in cuticle composition and thickening.

Genotyping of the *kdr* locus confirmed not only the presence, but probable fixation of the pyrethroid resistance mutation F1534C, in line with previous studies [23]. However, the V1016I pyrethroid resistant mutation showed a significant frequency increase (8% in 2009 [23], to 17% in 2013, this study; Fisher's exact tests, $P = 0.019$). The role of V1016I in resistance to pyrethroids is currently unclear but frequencies of both mutations appear informative and should be routinely monitored [50]. Hu et al [51] found that the F1534C mutation is more effective in reducing sensitivity of the sodium channel to type I than to type II pyrethroids. This could explain the higher resistance level of *Ae. aegypti* from Funchal to permethrin when compared to cyfluthrin. However, reversal of resistance to permethrin with the synergist PBO was comparable to that obtained for cyfluthrin suggesting that resistance to both type I and type II pyrethroids may be primarily mediated by the metabolic activity of cytochrome P450 oxidases. These findings should be taken into consideration by the Health Authorities in Madeira when deciding between type I or type II pyrethroids for vector control. If this chemical class remains an option for chemical control of *Ae. aegypti* in Madeira island, the use of type II pyrethroids in combination with the synergist PBO appears to be a more effective option.

Conclusion

Insecticide resistance mediated by multiple mechanisms was identified in *Ae. aegypti* from two localities in Madeira. In addition to target-site (*kdr*) and metabolic resistance, a third mechanism consisting of cuticle thickening may also be involved, confirming that the insecticide resistance phenotype is multifactorial, and consequently is likely to be challenging to reverse. The recent presence of this species in the island and the absence of a continuous, island-wide, insecticide-based control suggests that at least some, and maybe most, of the insecticide resistance mechanisms detected were already present in the colonizing specimens. Thus, the

resistance status of these mosquitoes may have played some role in the establishment of this vector in the island, despite the 2006–2008 insecticide-based vector control campaign. With the current knowledge regarding insecticide resistance status and identification of underlying mechanisms, resistance management strategies including mode of action rotation (such as bio-cides and insect growth regulators), as well as alternative to chemical-based vector control interventions (ranging from environmental management to new paradigms and biotechnol-ogy-based approaches) is strongly advised, to control *Ae. aegypti* and thus decrease the proba-bility of arbovirus transmission.

Supporting information

S1 Table. List of probes analysed after the quality control and primers used in qPCR.
(XLS)

Acknowledgments

We thank Patricia Pignatelli (Liverpool School of Tropical Medicine, UK) for her help in the microarray study.

Author Contributions

Conceptualization: Carla Alexandra Sousa.

Formal analysis: Gonalo Seixas, Linda Grigoraki, David Weetman, Joo Pinto, John Vontas, Carla Alexandra Sousa.

Funding acquisition: Joo Pinto, Carla Alexandra Sousa.

Investigation: Gonalo Seixas, Linda Grigoraki, David Weetman.

Methodology: Gonalo Seixas, Linda Grigoraki, David Weetman, Joo Pinto, John Vontas, Carla Alexandra Sousa.

Project administration: John Vontas, Carla Alexandra Sousa.

Resources: Gonalo Seixas, Linda Grigoraki, David Weetman, Ana Clara Silva, Joo Pinto, John Vontas, Carla Alexandra Sousa.

Supervision: Carla Alexandra Sousa.

Visualization: Gonalo Seixas, Linda Grigoraki, David Weetman, Jos Luis Vicente, Ana Clara Silva, Joo Pinto, John Vontas, Carla Alexandra Sousa.

Writing – original draft: Gonalo Seixas, Linda Grigoraki, David Weetman, Jos Luis Vicente, Ana Clara Silva, Joo Pinto, John Vontas, Carla Alexandra Sousa.

Writing – review & editing: Gonalo Seixas, Linda Grigoraki, David Weetman, Jos Luis Vicente, Ana Clara Silva, Joo Pinto, John Vontas, Carla Alexandra Sousa.

References

1. Kraemer MUG, Sinka ME, Duda KA, Mylne AQN, Shearer FM, Barker CM, et al. The global distribution of the arbovirus vectors *Aedes aegypti* and *Ae. albopictus*. *Elife*. 2015; 4: 1–18. <https://doi.org/10.7554/eLife.08347> PMID: 26126267
2. Beebe NW, Cooper RD, Mottram P, Sweeney AW. Australia's dengue risk driven by human adaptation to climate change. *PLoS Negl Trop Dis*. 2009; 3: e429. <https://doi.org/10.1371/journal.pntd.0000429> PMID: 19415109

3. Brown JE, Obas V, Morley V, Powell JR. Phylogeography and spatio-temporal genetic variation of *Aedes aegypti* (Diptera: Culicidae) populations in the Florida Keys. *J Med Entomol*. 2013; 50: 294–9. Available: <http://www.ncbi.nlm.nih.gov/pubmed/23540116> PMID: 23540116
4. Margarita Y, Santos Grácio AJ, Lencastre I, Silva AC, Novo T, Sousa C. First record of *Aedes* (Stegomyia) *aegypti* (Linnaeus, 1762) (Diptera, Culicidae) in Madeira Island—Portugal (Portuguese, English abstract). *Acta Parasitol Port*. 2006; 13: 59–61.
5. Almeida APG, Gonçalves YM, Novo MT, Sousa CA, Melim M, Gracio AJ. Vector monitoring of *Aedes aegypti* in the Autonomous Region of Madeira, Portugal. *Euro Surveill*. 2007; 12(46), pp.3311.
6. Gonçalves Y, Silva J, Biscoito M. On the presence of *Aedes* (Stegomyia) *aegypti* Linnaeus, 1762 (Insecta, Diptera, Culicidae) in the island of Madeira (Portugal). *Bol Mus Mun Funchal*. 2008; 58: 53–59.
7. European Centre for Disease Prevention and Control. Dengue outbreak in Madeira, Portugal, March 2013. Stockholm: ECDC; 2014.
8. Gould EA, Gallian P, De Lamballerie X, Charrel RN. First cases of autochthonous dengue fever and chikungunya fever in France: from bad dream to reality! *Clin Microbiol Infect*. Blackwell Publishing Ltd; 2010; 16: 1702–1704. <https://doi.org/10.1111/j.1469-0691.2010.03386.x> PMID: 21040155
9. Gjenero-Margan I, Aleraj B, Krajcar D, Lesnikar V, Klobučar a, Pem-Novosel I, et al. Autochthonous dengue fever in Croatia, August–September 2010. *Euro Surveill*. 2011; 16. Available: <http://www.ncbi.nlm.nih.gov/pubmed/21392489>
10. Succo T, Leparco-Goffart I, Ferré J, Roiz D, Broche B, Maquart M, et al. Autochthonous dengue outbreak in Nîmes, South of France, July to September 2015. *Eurosurveillance*. 2016; 21: 1–7. <https://doi.org/10.2807/1560-7917.ES.2016.21.21.30240> PMID: 27254729
11. Vontas J, Kioulos E, Pavlidi N, Morou E, Torre A, Ranson H. Insecticide resistance in the major dengue vectors *Aedes albopictus* and *Aedes aegypti*. *Pestic Biochem Physiol*. Elsevier Inc.; 2012; 6–11. <https://doi.org/10.1016/j.pestbp.2012.05.008>
12. Smith LB, Kasai S, Scott JG. Pyrethroid resistance in *Aedes aegypti* and *Aedes albopictus*: Important mosquito vectors of human diseases. *Pestic Biochem Physiol*. Elsevier B.V.; 2016; <https://doi.org/10.1016/j.pestbp.2016.03.005> PMID: 27742355
13. Stevenson BJ, Pignatelli P, Nikou D, Paine MJL. Pinpointing P450s associated with pyrethroid metabolism in the dengue vector, *Aedes aegypti*: developing new tools to combat insecticide resistance. *PLoS Negl Trop Dis*. 2012; 6: e1595. <https://doi.org/10.1371/journal.pntd.0001595> PMID: 22479665
14. Grigoraki L, Balabanidou V, Meristoudis C, Miridakis A, Ranson H, Swevers L, et al. Functional and immunohistochemical characterization of CCEae3a, a carboxylesterase associated with temephos resistance in the major arbovirus vectors *Aedes aegypti* and *Ae. albopictus*. *Insect Biochem Mol Biol*. 2016; 74: 61–67. <https://doi.org/10.1016/j.ibmb.2016.05.007> PMID: 27180726
15. Nkya TE, Akhouayri I, Kisinza W, David JP. Impact of environment on mosquito response to pyrethroid insecticides: Facts, evidences and prospects. *Insect Biochem Mol Biol*. Elsevier Ltd; 2012; 43: 407–416. <https://doi.org/10.1016/j.ibmb.2012.10.006> PMID: 23123179
16. Pedrini N, Mijailovsky SJ, Girotti JR, Stariolo R, Cardozo RM, Gentile A, et al. Control of pyrethroid-resistant chagas disease vectors with entomopathogenic fungi. *PLoS Negl Trop Dis*. 2009; 3. <https://doi.org/10.1371/journal.pntd.0000434> PMID: 19434231
17. Wood OR, Hanrahan S, Coetzee M, Koekemoer LL, Brooke BD. Cuticle thickening associated with pyrethroid resistance in the major malaria vector *Anopheles funestus*. *Parasit Vectors*. 2010; 3: 67. Available: <http://www.parasitesandvectors.com/content/3/1/67> <https://doi.org/10.1186/1756-3305-3-67> PMID: 20684757
18. Balabanidou V, Kampouraki A, MacLean M, Blomquist GJ, Tittiger C, Juárez MP, et al. Cytochrome P450 associated with insecticide resistance catalyzes cuticular hydrocarbon production in *Anopheles gambiae*. *Proc Natl Acad Sci*. 2016; 113: 201608295. <https://doi.org/10.1073/pnas.1608295113> PMID: 27439866
19. Vontas J, David JP, Nikou D, Hemingway J, Christophides GK, Louis C, et al. Transcriptional analysis of insecticide resistance in *Anopheles stephensi* using cross-species microarray hybridization. *Insect Mol Biol*. 2007; 16: 315–324. <https://doi.org/10.1111/j.1365-2583.2007.00728.x> PMID: 17433071
20. David J-P, Coissac E, Melodelima C, Poupardin R, Riaz MA, Chandor-Proust A, et al. Transcriptome response to pollutants and insecticides in the dengue vector *Aedes aegypti* using next-generation sequencing technology. *BMC Genomics*. 2010; 11: 216. <https://doi.org/10.1186/1471-2164-11-216> PMID: 20356352
21. Fang F, Wang W, Zhang D, Lv Y, Zhou D, Ma L, et al. The cuticle proteins: a putative role for deltamethrin resistance in *Culex pipiens pallens*. *Parasitol Res*. 2015; 114: 4421–4429. <https://doi.org/10.1007/s00436-015-4683-9> PMID: 26337265

22. Ishak IH, Riveron JM, Ibrahim SS, Stott R, Longbottom J, Irving H, et al. The Cytochrome P450 gene CYP6P12 confers pyrethroid resistance in *kdr*-free Malaysian populations of the dengue vector *Aedes albopictus*. *Sci Rep. Nature Publishing Group*; 2016; 6: 24707. <https://doi.org/10.1038/srep24707> PMID: 27094778
23. Seixas G, Salgueiro P, Silva AC, Campos M, Spenassatto C, Reyes-lugo M, et al. *Aedes aegypti* on Madeira Island (Portugal): genetic variation of a recently introduced dengue vector. 2013; 1–8. <https://doi.org/10.1590/0074-0276130386> PMID: 24473797
24. Nazareth T, Teodósio R, Porto G, Gonçalves L, Seixas G, Silva AC, et al. Strengthening the perception-assessment tools for dengue prevention: a cross-sectional survey in a temperate region (Madeira, Portugal). *BMC Public Health*. 2014; 14: 39. <https://doi.org/10.1186/1471-2458-14-39> PMID: 24428823
25. Oliveira P, Pereira PT. Who values what in a tourism destination? The case of Madeira Island. *Tour Econ*. 2008; 14: 155–168. <https://doi.org/10.5367/000000008783554758>
26. World Health Organization. Test procedures for insecticide resistance monitoring in malaria vector mosquitoes. Geneva: World Health Organization; 2013.
27. World Health Organization. Monitoring and managing insecticide resistance in *Aedes* mosquito populations Interim guidance for entomologists. Geneva: World Health Organization; 2016.
28. World Health Organization. Techniques to detect insecticide resistance mechanisms field and laboratory manual. Geneva: World Health Organization; 1998.
29. Moran MD. Arguments for rejecting the sequential Bonferroni in ecological studies. *Oikos*. 2003; 100: 403–405. <https://doi.org/10.1034/j.1600-0706.2003.12010.x>
30. Poupardin R, Srisukontarat W, Yunta C, Ranson H. Identification of carboxylesterase genes implicated in temephos resistance in the dengue vector *Aedes aegypti*. *PLoS Negl Trop Dis*. 2014; 8. <https://doi.org/10.1371/journal.pntd.0002743> PMID: 24651719
31. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res*. 2001 May 1; 29:e45 PMID: 11328886
32. Collins FH, Mendez M a, Rasmussen MO, Mehaffey PC, Besansky NJ, Finnerty V. A ribosomal RNA gene probe differentiates member species of the *Anopheles gambiae* complex. *Am J Trop Med Hyg*. 1987; 37: 37–41. Available: <http://www.ncbi.nlm.nih.gov/pubmed/2886070> PMID: 2886070
33. Saavedra-Rodriguez K, Urdaneta-Marquez L, Rajatileka S, Moulton M, Flores a E, Fernandez-Salas I, et al. A mutation in the voltage-gated sodium channel gene associated with pyrethroid resistance in Latin American *Aedes aegypti*. *Insect Mol Biol*. 2007; 16: 785–98. <https://doi.org/10.1111/j.1365-2583.2007.00774.x> PMID: 18093007
34. Harris AF, Rajatileka S, Ranson H. Pyrethroid resistance in *Aedes aegypti* from Grand Cayman. *Am J Trop Med Hyg*. 2010; 83: 277–84. <https://doi.org/10.4269/ajtmh.2010.09-0623> PMID: 20682868
35. David J-P, Faucon F, Chandor-Proust A, Poupardin R, Riaz MA, Bonin A, et al. Comparative analysis of response to selection with three insecticides in the dengue mosquito *Aedes aegypti* using mRNA sequencing. *BMC Genomics*. 2014; 15: 174. <https://doi.org/10.1186/1471-2164-15-174> PMID: 24593293
36. Bingham G, Strode C, Tran L, Khoa PT, Jamet HP. Can piperonyl butoxide enhance the efficacy of pyrethroids against pyrethroid-resistant *Aedes aegypti*? *Trop Med Int Health*. 2011; 16: 492–500. <https://doi.org/10.1111/j.1365-3156.2010.02717.x> PMID: 21324051
37. Coleman M, Hemingway J. Insecticide resistance monitoring and evaluation in disease transmitting mosquitoes. *J Pestic Sci*. 2007; 32: 69–76. <https://doi.org/10.1584/jpestics.R07-01>
38. Morou E, Dowd AJ, Rajatileka S, Steven A, Hemingway J, Ranson H, et al. A simple colorimetric assay for specific detection of glutathione-S transferase activity associated with DDT resistance in mosquitoes. *PLoS Negl Trop Dis*. 2010; 4: 1–6. <https://doi.org/10.1371/journal.pntd.0000808> PMID: 20824165
39. Strode C, Wondji CS, David J, Hawkes NJ, Lumjuan N, Nelson DR, et al. Genomic analysis of detoxification genes in the mosquito *Aedes aegypti*. *Insect Biochem Mol Biol*. 2008; 38: 113–123. <https://doi.org/10.1016/j.ibmb.2007.09.007> PMID: 18070670
40. Pavlidi N, Monastiriotti M, Daborn P, Livadaras I, Van Leeuwen T, Vontas J. Transgenic expression of the *Aedes aegypti* CYP9J28 confers pyrethroid resistance in *Drosophila melanogaster*. *Pestic Biochem Physiol*. Elsevier Inc.; 2012; 104: 132–135. <https://doi.org/10.1016/j.pestbp.2012.07.003>
41. Grigoraki L, Lagnel J, Kioulos I, Kampouraki A, Morou E, Labbé P, et al. Transcriptome profiling and genetic study reveal amplified carboxylesterase genes implicated in temephos resistance, in the asian tiger mosquito *Aedes albopictus*. *PLoS Negl Trop Dis*. 2015; 9: e0003771. <https://doi.org/10.1371/journal.pntd.0003771> PMID: 26000638
42. Kasai S, Komagata O, Itokawa K, Shono T, Ng LC, Kobayashi M, et al. Mechanisms of pyrethroid resistance in the dengue mosquito vector, *Aedes aegypti*: target site Insensitivity, penetration, and metabolism. *PLoS Negl Trop Dis*. 2014; 8. <https://doi.org/10.1371/journal.pntd.0002948> PMID: 24945250

43. Ahmad M, Denholm I, Bromilow RH. Delayed cuticular penetration and enhanced metabolism of delta-methrin in pyrethroid-resistant strains of *Helicoverpa armigera* from China and Pakistan. *Pest manag Sci*. 2006; 62: 805–810. <https://doi.org/10.1002/ps.1225> PMID: 16649192
44. Strycharz JP, Lao A, Li H, Qiu X, Lee SH, Sun W, et al. Resistance in the highly DDT-resistant 91-R strain of *Drosophila melanogaster* involves decreased penetration, increased metabolism, and direct excretion. *Pestic Biochem Physiol*. Elsevier Inc.; 2013; 107: 207–217. <https://doi.org/10.1016/j.pestbp.2013.06.010>
45. Burmester T. Evolution and function of the insect hexamerins. *Eur J Entomol*. 1999; 96: 213–225.
46. Chrysanthis G, Kaliafas AD, Mintzas AC. Biosynthesis and tissue distribution of four major larval serum proteins during development of *Ceratitis capitata* (Diptera). *Insect Biochem Mol Biol*. 1994; 24: 811–818. [https://doi.org/10.1016/0965-1748\(94\)90109-0](https://doi.org/10.1016/0965-1748(94)90109-0)
47. Gefen E, Talal S, Brendzel O, Dror A, Fishman A. Variation in quantity and composition of cuticular hydrocarbons in the scorpion *Buthus occitanus* (Buthidae) in response to acute exposure to desiccation stress. *Comp Biochem Physiol -Part A Mol Integr Physiol*. Elsevier Inc.; 2015; 182: 58–63. <https://doi.org/10.1016/j.cbpa.2014.12.004> PMID: 25499238
48. Fedorka KM, Copeland EK, Winterhalter WE. Seasonality influences cuticle melanization and immune defense in a cricket: support for a temperature-dependent immune investment hypothesis in insects. *J Exp Biol*. 2013; 216: 4005–10. <https://doi.org/10.1242/jeb.091538> PMID: 23868839
49. Arcaz AC, Huestis DL, Dao A, Yaro AS, Diallo M, Andersen J, et al. Desiccation tolerance in *Anopheles coluzzii*: the effects of spiracle size and cuticular hydrocarbons. *J Exp Biol*. 2016; 219: 1675–1688. <https://doi.org/10.1242/jeb.135665> PMID: 27207644
50. Vera-Maloof FZ, Saavedra-Rodriguez K, Elizondo-Quiroga AE, Lozano-Fuentes S, Black IV WC. Coevolution of the Ile1,016 and Cys1,534 mutations in the voltage gated sodium channel gene of *Aedes aegypti* in Mexico. *PLoS Negl Trop Dis*. 2015; 9: 1–22. <https://doi.org/10.1371/journal.pntd.0004263> PMID: 26658798
51. Hu Z, Du Y, Nomura Y, Dong K. A sodium channel mutation identified in *Aedes aegypti* selectively reduces cockroach sodium channel sensitivity to type I, but not type II pyrethroids. *Insect Biochem Mol Biol*. Elsevier Ltd; 2011; 41: 9–13. <https://doi.org/10.1016/j.ibmb.2010.09.005> PMID: 20869441