



ELSEVIER

Contents lists available at ScienceDirect

## European Journal of Obstetrics & Gynecology and Reproductive Biology

journal homepage: [www.elsevier.com/locate/ejogrb](http://www.elsevier.com/locate/ejogrb)



### Review article



## EBCOG position statement on COVID-19 vaccination for pregnant and breastfeeding women

Inês Martins<sup>a</sup>, Frank Louwen<sup>b</sup>, Diogo Ayres-de- Campos<sup>a,c</sup>, Tahir Mahmood<sup>d,\*</sup>

<sup>a</sup> Department of Obstetrics and Gynecology, Santa Maria University Hospital, Lisbon, Portugal

<sup>b</sup> Goethe-Universität, Frankfurt am Main, Germany

<sup>c</sup> Medical School, University of Lisbon, Portugal

<sup>d</sup> EBCOG Standards of Care and Position Statement Group, Victoria Hospital, Kirkcaldy, United Kingdom

#### ARTICLE INFO

##### Article history:

Received 4 May 2021

Accepted 11 May 2021

Available online xxx

##### Keywords:

Covid-19

Pregnancy

Vaccination

EBCOG

Position statement

Breast feeding

Pre-term labour

Maternal mortality

#### ABSTRACT

Covid 19 pandemic has led to significant mortality and long term morbidity globally. Pregnant women are at increased risk of severe illness from COVID 19 infection. There is an urgent need for all health authorities and Governments to offer vaccination to all pregnant women especially those with high risk pregnancy.

© 2021 Elsevier B.V. All rights reserved.

\* Corresponding author.

E-mail address: [tahir.mahmood@nhs.scot](mailto:tahir.mahmood@nhs.scot) (T. Mahmood).

<https://doi.org/10.1016/j.ejogrb.2021.05.021>

0301-2115/© 2021 Elsevier B.V. All rights reserved.

Contents

The risks of Covid-19 infection during pregnancy and breastfeeding ..... 00  
Covid-19 vaccines ..... 00  
EBCOG position ..... 00  
EBCOG supports that COVID-19 vaccination be recommended to all breastfeeding women, in the absence of a specific contraindication ... 00  
References ..... 00

**The risks of Covid-19 infection during pregnancy and breastfeeding**

There is now strong evidence that pregnant women are at increased risk for severe illness from COVID-19 when compared with non-pregnant women, particularly when infection occurs during the third trimester of pregnancy [1–3]. The risk of intensive care unit admission is around 1% in this population, and the risk of invasive mechanical ventilation around 0.3% [1,2]. These values are *circa* three times higher than those of the non-pregnant population of the same age. While the absolute risk of death from COVID-19 in pregnant women remains low, at about 0.15% in symptomatic patients, it is around 70% higher than that of non-pregnant women of the same age [1,2]. Maternal conditions such as age over 40 years, obesity, chronic hypertension, and preexisting diabetes are additional risk factors for severe disease [1,2,4].

There is also evidence that COVID-19 infection during pregnancy increases the risk of preterm birth about three-fold, mainly due to iatrogenic preterm delivery [1,2]. Vertical transmission of SARS-CoV-2 [5] occurs in approximately 2.5% of pregnant women and appears to be mainly limited to cases of third-trimester infection [6].

For all the reasons described above, pregnant women need to be considered a high-risk population for serious COVID-19 infection, and there are clear benefits for both mother and fetus of avoiding this disease during pregnancy [7,8]. Breastfeeding women are believed to have similar risks of serious disease as non-pregnant women.

**Covid-19 vaccines**

Currently authorized COVID-19 vaccines in Europe use different technologies, including messenger RNA (mRNA), viral vectors, and recombinant proteins produced in a baculovirus system. However, none of them use live viruses, therefore they are not capable of causing disease [9,10]. Other vaccines that do not use live viruses show no evidence of increased risk during human pregnancy and breastfeeding [11].

Existing vaccines do not interact with individuals' DNA or cause genetic mutations [9–11]. Animal studies did not reveal any safety concerns regarding female reproduction, embryo-fetal or neonatal development [9,11].

All these vaccines were rigorously evaluated in randomised controlled trials, but pregnancy and breastfeeding were exclusion criteria [9–11]. There is limited data from pregnant women who were inadvertently enrolled in these trials, but it is reassuring on the safety of vaccination in pregnancy [9,11]. Self-reported information after ongoing COVID-19 vaccination in the USA (86 956 pregnant women, as of the 12th April 2021 [12], of which 35 691 have been the subject of a publication [13]) has not revealed any safety issues, but follow-up is naturally short. It is strongly believed that COVID-19 vaccines do not cause any harm to newborns by way of breastmilk [14].

In a small prospective study [15] comparing 84 pregnant, 31 breastfeeding, and 16 non-pregnant women vaccinated with a mRNA vaccine, vaccine-induced immune responses were similar,

and higher antibody titers were found after vaccination than after infection in pregnancy. Vaccine-generated antibodies were found in umbilical cord blood and breast milk samples, showing a potential for fetal and neonatal immunization with maternal vaccination.

Of late, several cases of cerebral vessels thrombotic (CVT) events following immunization with COVID-19 viral vector vaccines (Astra Zeneca and Johnson & Johnson) have been reported, more commonly in women. Nevertheless, they remain a rare event, with an incidence of about 4.1 per million compared with the incidence of 39.0 per million people with a diagnosis of Covid [16]. The European Medicine Agency (EMA) and the United Kingdom Medicines & Healthcare products Regulatory Agency (MHRA) have stated that the benefits of vaccination clearly outweigh the risks. Currently there is no evidence on the risk of thrombotic events following immunization with COVID-19 viral vector vaccines in pregnancy, a well-known prothrombotic state. Women should be counseled as regards this potential risk, and healthcare professionals should have a low threshold for investigation, if a thrombotic event is suspected [17,18].

**EBCOG position**

All vaccination during pregnancy is naturally optional, but healthcare providers need to advise women about the risks of acquiring COVID-19 infection in pregnancy, as well as the limited existing evidence on the benefits and potential side effects of existing vaccines, as a basis to shared decision-making on this issue. EBCOG acknowledges that a clear indication from healthcare providers would improve adherence to vaccination programs and reduce women's anxiety with the decision [7], but this is not possible with the currently existing evidence.

EBCOG acknowledges that there is limited evidence on the long-term safety of COVID-19 vaccination during pregnancy, and that there is a need for more robust data before it can be recommended to all pregnant women. Nevertheless, **the possibility of vaccination should be offered to all pregnant women, after being adequately informed of the benefits and risks. EBCOG urges all Health Authorities and Governments to make vaccination available to all pregnant women wishing to take them.**

In women with the co-morbidities listed above, and in those at higher-risk of exposure to the disease, EBCOG considers that the benefits are likely to outweigh the predictable risks, and therefore **vaccination should be recommended to high-risk pregnant women, in the absence of contraindications.** There is currently insufficient data regarding the preferential use of a particular COVID-19 vaccine.

**EBCOG supports that COVID-19 vaccination be recommended to all breastfeeding women, in the absence of a specific contraindication**

The position statement has been approved by the officers group of EBCOG and the executive board of EBCOG.

## References

- [1] Zambrano LD, Ellington S, Strid P, et al. Update: characteristics of symptomatic women of reproductive age with laboratory-confirmed SARS-CoV-2 infection by pregnancy status - United States, January 22–October 3, 2020. CDC COVID-19 Response Pregnancy and Infant Linked Outcomes Team. *MMWR Morb Mortal Wkly Rep* 2020;69:1641–7.
- [2] Allotey J, Stallings E, Bonet M, et al. Clinical manifestations, risk factors, and maternal and perinatal outcomes of coronavirus disease 2019 in pregnancy: living systematic review and meta-analysis. *BMJ* 2020;370:m3320.
- [3] Delahoy MJ, Whitaker M, O'Halloran A, et al. Characteristics and maternal and birth outcomes of hospitalized pregnant women with laboratory-confirmed COVID-19 - COVID-NET, 13 states, March 1–August 22, 2020. COVID-NET Surveillance Team. *MMWR Morb Mortal Wkly Rep* 2020;69:1347–54.
- [4] Panagiotakopoulos L, Myers TR, Gee J, et al. SARS-CoV-2 infection among hospitalized pregnant women: reasons for admission and pregnancy characteristics - eight U.S. health care centers, March 1–May 30, 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:1355–9.
- [5] Blumberg DA, Underwood MA, Hedriana HL, Lakshminrusimha S. Vertical transmission of SARS-CoV-2: what is the optimal definition? *Am J Perinatol* 2020;37(June(8)):769–72.
- [6] Goh XL, Low YF, Ng CH, Amin Z, Ng YPM. Incidence of SARS-CoV-2 vertical transmission: a meta-analysis. *Arch Dis Child Fetal Neonatal Ed* 2021;106(January(1)):112–3.
- [7] Chervenak FA, McCullough LB, Bornstein E, Johnson L, Katz A, et al. Professionally responsible coronavirus disease 2019 vaccination counseling of obstetrical and gynecologic patients. *Am J Obstet Gynecol* 2021(February) S0002-9378(21)00082-X.
- [8] Nassar AH, Visser GHA, Nicholson WK, et al. FIGO safe motherhood, newborn health committee. FIGO statement: vaccination in pregnancy. *Int J Gynaecol Obstet* 2021;152(February(2)):139–43.
- [9] European Medicines Agency. Coronavirus disease (COVID-19). Available at: <https://www.ema.europa.eu/en/human-regulatory/overview/public-health-threats/coronavirus-disease-covid-19>.
- [10] Rasmussen SA, Kelley CF, Horton JP, et al. Coronavirus Disease 2019 (COVID-19) Vaccines and pregnancy: what obstetricians need to know. *Obstet Gynecol* 2021;137(March (3)):408–14.
- [11] Centers for Disease Control and Prevention. Information about COVID-19 Vaccines for People who Are Pregnant or Breastfeeding. Available at: <https://www.cdc.gov/coronavirus/2019-nCoV/vaccines/recommendations/pregnancy.html>.
- [12] Centers for Disease Control and Prevention. V-safe COVID-19 Vaccine Pregnancy Registry. Available at: <https://www.cdc.gov/coronavirus/2019-nCoV/vaccines/safety/vsafepregnancyregistry.html>.
- [13] Shimabukuro TT, Kim SY, Myers TR, et al. Preliminary findings of mRNA Covid-19 vaccine safety in pregnant persons. *New Eng J Med* 2021 published online April 21 2021.
- [14] Dooling K, Marin M, Wallace M, et al. The Advisory Committee on Immunization Practices' updated interim recommendation for allocation of COVID-19 vaccine - United States, December 2020. *MMWR Morb Mortal Wkly Rep* 2021;69:1657–60.
- [15] Gray KJ, Bordt EA, Atyeo C, Deriso E, Akinwunmi B, Young N, et al. COVID-19 vaccine response in pregnant and lactating women: a cohort study. *medRxiv* 2021(March) [Preprint] 2021.03.07.21253094.
- [16] Torjesen Ingrid. Risk of cerebral blood clots from Covid 19 is 10 times that from vaccination, study finds. *BMJ* 2021;373: n1005.
- [17] British Society for Haematology. Guidance produced by the expert haematology panel (EHP) focused on vaccine induced thrombosis and thrombocytopenia (VITT). Available at: 2021. <https://b-s-h.org.uk/>.
- [18] The Royal College of Emergency Medicine. The society for acute medicine & the royal college of physicians joint guidance. Available at: 2021. <https://www.rcem.ac.uk/docs/Policy/ED-M%20Vaccine%20pathway%20concerns%20-%20RCP%20-%20SAM%20-%20RCEM.pdf>.