

Cite as: K. Bjornevik *et al.*, *Science* 10.1126/science.abj8222 (2022).

# Longitudinal analysis reveals high prevalence of Epstein-Barr virus associated with multiple sclerosis

Kjetil Bjornevik<sup>1,†</sup>, Marianna Cortese<sup>1,†</sup>, Brian C. Healy<sup>2,3,4</sup>, Jens Kuhle<sup>5</sup>, Michael J. Mina<sup>6,7,8</sup>, Yumei Leng<sup>6</sup>, Stephen J. Elledge<sup>6</sup>, David W. Niebuhr<sup>9</sup>, Ann I. Scher<sup>9</sup>, Kassandra L. Munger<sup>1,10,11,‡\*</sup>, Alberto Ascherio<sup>1,10,11,‡\*</sup>

<sup>1</sup>Department of Nutrition, Harvard T. H. Chan School of Public Health, Boston, MA, USA. <sup>2</sup>Partners Multiple Sclerosis Center, Brigham and Women's Hospital, Boston, MA, USA. <sup>3</sup>Department of Neurology, Harvard Medical School, Boston, MA, USA. <sup>4</sup>Biostatistics Center, Massachusetts General Hospital, Boston, MA, USA. <sup>5</sup>Neurologic Clinic and Polyclinic, MS Center and Research Center for Clinical Neuroimmunology and Neuroscience Basel (RC2NB), University Hospital Basel, University of Basel, Basel, Switzerland. <sup>6</sup>Division of Genetics, Brigham and Women's Hospital, Howard Hughes Medical Institute, Department of Genetics, and Program in Virology, Harvard Medical School, Boston, MA, USA. <sup>7</sup>Center for Communicable Disease Dynamics, Department of Epidemiology, and Department of Immunology and Infectious Diseases, Harvard T. H. Chan School of Public Health, Boston, MA, USA. <sup>8</sup>Department of Pathology, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA. <sup>9</sup>Department of Preventive Medicine and Biostatistics, Uniformed Services University of the Health Sciences, Bethesda, MD, USA. <sup>10</sup>Department of Epidemiology, Harvard T. H. Chan School of Public Health, Boston, MA, USA. <sup>11</sup>Channing Laboratory, Department of Medicine, Brigham and Women's Hospital, and Harvard Medical School, Boston, MA, USA.

†These authors contributed equally to this work.

‡These authors contributed equally to this work.

\*Corresponding author. Email: [aascheri@hsph.harvard.edu](mailto:aascheri@hsph.harvard.edu)

**Multiple sclerosis (MS) is a chronic inflammatory demyelinating disease of the central nervous system of unknown etiology. We tested the hypothesis that MS is caused by Epstein-Barr virus (EBV) in a cohort comprising more than 10 million young adults on active duty in the US military, 955 of whom were diagnosed with MS during their period of service. Risk of MS increased 32-fold after infection with EBV but was not increased after infection with other viruses, including the similarly transmitted cytomegalovirus. Serum levels of neurofilament light chain, a biomarker of neuroaxonal degeneration, increased only after EBV seroconversion. These findings cannot be explained by any known risk factor for MS and suggest EBV as the leading cause of MS.**

Multiple sclerosis (MS) is a chronic inflammatory demyelinating disease of the central nervous system of unknown etiology. The demyelination in the brain and spinal cord is an immune-mediated process (1) possibly triggered by a viral infection (2). Among the putative causal agents, the top candidate is Epstein-Barr virus (EBV) (3). EBV is a human herpesvirus that after infection persists in latent form in B lymphocytes throughout the life of the host (3). A causal role of EBV is supported by the increased MS risk after infectious mononucleosis (4), elevated serum antibody titers against EBV nuclear antigens (EBNAs) (5), and by the presence of EBV in MS demyelinated lesions reported in some (6–8), but not all (9), pathological studies. Evidence of causality, however, remains inconclusive.

Causality implies that some individuals who developed MS after EBV infection would not have developed MS if they had not been infected with EBV. Ruling out a randomized trial, the gold standard to study this counterfactual occurrence is an “experiment of nature,” a longitudinal investigation of MS incidence in a cohort of EBV-negative individuals, some of whom will be infected with EBV during the follow-up and some who will not. The ubiquitous nature of EBV, which infects ~95% of adults, and the fact that MS is a relatively rare disease, has until now impeded such an

investigation. Over the course of a 20-year collaboration with the US military, we have identified cases of MS in a cohort composed of active-duty US military personnel between 1993 and 2013, a racially diverse population of >10 million individuals. Active-duty members are screened for HIV at the start of military service and biennially thereafter, and residual serum from these tests (>62 million serum samples) is archived in the Department of Defense Serum Repository (DoDSR) (10). We used samples stored in the DoDSR to determine EBV status at time of first sample and the relation between EBV infection and MS onset during the period of active duty. In a preliminary study, we found that 5.3% of individuals were EBV-negative at the time of first sample (11), corresponding to hundreds of thousands of EBV-negative young adults at risk of EBV infection and MS.

We documented 955 incident MS cases among active-duty military personnel [including 315 cases from our preliminary study (11)]. For each MS case, we identified up to three serum samples collected before the date of MS onset (the first available, the last collected before disease onset, and one in between). Cases were matched to two randomly selected individuals without MS of the same age, sex, race/ethnicity, branch of military service, and dates of collection of blood samples who were on active military duty when the case was

diagnosed (Fig. 1A and fig. S1). There were 801 MS cases and 1566 controls with samples available to assess EBV infection status. Most of the individuals in our study were <20 years of age at the time of their first blood collection (Fig. 1B), and those who developed MS had symptom onset a median of 10 years after time of first sample (Fig. 1C).

Only one of the 801 MS cases occurred in an individual who was EBV-negative in the last sample, which was collected at a median of 1 year before MS onset [hazard ratio (HR) for MS comparing EBV-positive versus EBV-negative = 26.5; 95% confidence interval (CI): 3.7 to 191.6;  $P = 0.001$ , conditional logistic regression]. At baseline, 35 MS cases and 107 controls were EBV-negative. All but one of these 35 EBV-negative MS cases became infected with EBV during the follow-up, and all seroconverted before the onset of MS (fig. S3). The median time from the first EBV-positive sample to MS onset was 5 years (range: 0 to 10 years), and the median time from estimated EBV seroconversion, defined as the midpoint between the last seronegative sample and the first seropositive sample, to MS onset was 7.5 years (range: 2 to 15 years). The high seroconversion rate among individuals who developed MS during follow-up (97%) contrasts with the 57% rate of seroconversion observed among individuals who did not develop MS (Fig. 2A), a rate consistent with previous reports among EBV-negative young adults (12). The HR for MS comparing EBV seroconversion versus persistent EBV seronegativity was 32.4 (95% CI: 4.3 to 245.3,  $P < 0.001$ ) (Fig. 2C).

Behavioral, environmental, or personal characteristics may correlate with a predisposition to both infection and MS. To assess this possibility, we measured antibodies against cytomegalovirus (CMV), a herpesvirus that, like EBV, is transmitted through the saliva. CMV displays socioeconomic and racial/ethnic disparities in age at infection in the US population (13) similar to those of EBV (14), thus constituting an ideal negative control (15). Among those who were CMV-negative at baseline, seroconversion for CMV occurred at a similar rate in those who later developed MS and those who did not (Fig. 2B). MS risk was lower among CMV-positive than among CMV-negative individuals (Fig. 2D), consistent with a previous report and with suggestions that the immune response to CMV attenuates the adverse effects of EBV (16).

Similar to other neurological diseases, the pathological mechanisms underlying MS likely start several years before the first symptoms appear (17). To further elucidate the temporal relation between EBV infection and MS, we measured serum concentrations of neurofilament light chain (sNfL), a sensitive, albeit not disease-specific, biomarker of ongoing neuroaxonal degeneration (18), using an ultrasensitive single-molecule assay (19) in the samples from those who were EBV-negative at baseline. We have previously reported that sNfL levels increase as early as 6 years before clinical MS onset and may be a more accurate marker of the time of initiation of

the disease process (20). sNfL levels in individuals who were EBV-negative at baseline and went on to develop MS were similar to those of non-MS controls before and around the time of EBV infection but increased after EBV infection (Fig. 3, A to C, and fig. S4). Thus, there were no signs of neuroaxonal degeneration before EBV seroconversion in individuals who later developed MS, indicating that EBV infection preceded not only symptom onset but also the time of the first detectable pathological mechanisms underlying MS.

To further explore whether immune dysregulation during the preclinical phase could increase susceptibility to viral infections more generally, we randomly selected 30 MS cases and 30 matched controls with serum samples collected shortly before (median: 1 year, range: 0 to 3) and soon after symptom onset (median: 1 year, range: 0 to 2) and conducted a comprehensive agnostic search of the anti-virome antibody response using VirScan, an assay based on a T7 phage-display immunoprecipitation and sequencing technology (21), which encodes the full proteome of most known human pathogenic viruses (~200 species, ~110,000 peptides) in 56-amino acid peptides with 28-amino acid overlap between adjacent peptides. VirScan thus enables comprehensive unbiased detection of antibodies raised against all linear peptides encoded in the genomes of viruses known to infect humans. The overall antibody response to viral peptides was similar in cases and controls at both time points, except for EBV (Fig. 4, A to D), arguing that the preclinical and early clinical phases in MS are not associated with immune dysregulation affecting general susceptibility to infections. Using a Z score of  $>3.5$  as an enrichment cutoff for identifying the presence of a peptide-specific antibody (22), the number of antibody-recognized EBV peptides with a nominally significant difference between MS cases and controls clearly stood out, both in the pre-onset (Fig. 4, E and G, and table S1) and post-onset (Fig. 4, F and H, and table S2) samples, which supports the specificity of the association between EBV and MS and argues against a second hit from another virus playing a major role in MS etiology.

A causal interpretation of our results requires ruling out the possibility that systematic differences between individuals who seroconverted and those who remained EBV-negative explain the results. These differences can be grouped into two categories: (i) confounding by known or unknown factors and (ii) reverse causation.

Confounding by known factors is virtually ruled out by the strength of the association. To explain a 32-fold increase in MS risk, any confounder would have to confer a >60-fold increase in risk of EBV seroconversion and a >60-fold risk of MS (23). None of the known or suspected risk factors for MS has such strong associations. The next strongest known risk factor for MS, homozygosity for the HLA-DR15 allele, which confers a threefold increase in MS risk (24), is not associated

with EBV positivity (25) and thus cannot explain the EBV-MS association. Rather, there is epidemiological (26) and experimental (27) evidence that EBV infection and HLA-DR15 may act synergistically in causing MS. Environmental factors are also far too weak to materially confound the EBV-MS association (28). The existence of a still unknown factor that increases the risk of both EBV infection and MS by >60-fold is rather implausible and there are no good candidates, even hypothetical ones. This conclusion would be robust even in the very unlikely case that EBV seroconversion in one of the MS cases was a false-positive result, in which case EBV infection would confer a 16-fold increase in MS risk.

Reverse causation could occur if the immune dysregulation during the preclinical phase of MS increases the susceptibility to EBV infection. In our agnostic search of the entire human virome during the preclinical phase of MS, we did not find other systematic differences in the antibody response to any pathogen except EBV that was related to previous infections in MS cases and controls, which makes it unlikely that immune dysregulation during this phase increases susceptibility to infections. This is consistent with previous studies reporting no difference in the frequency of infections in the 5 years preceding MS onset (29) or in individuals with untreated MS (30). Although in one study, hospitalizations for bacterial infections in adolescence were associated with MS risk, this association was modest and therefore cannot explain our study results (31). Additional arguments against reverse causality are that EBV seroconversion occurs before elevation of sNfL levels, which is an early marker of preclinical MS, and the long lag time (median: 7.5 years) between EBV infection and MS clinical onset. The increased MS risk 15 years or longer after infectious mononucleosis (32) and the observation that anti-EBNA antibodies are a strong and consistent predictor of MS risk in EBV-positive individuals up to 15 to 20 years later (33) provide further and independent evidence against reverse causation. Collectively, these findings strongly suggest that the occurrence of EBV infection, detectable by the elicited immune response, is a cause and not a consequence of MS.

One MS case was EBV-negative in the last sample, obtained 3 months before MS onset, which could suggest that EBV was not the cause of disease in this patient. This individual could have been infected with EBV after the last blood collection, could have failed to seroconvert in response to infection (an uncommon but nevertheless regular phenomenon seen after infections and vaccines), or could have been misdiagnosed. Another explanation is related to etiological diversity, which is common for any clinically defined disease. For example, all cases of paralytic poliomyelitis are by definition caused by poliovirus, but rare cases of acute flaccid paralysis, clinically indistinguishable from poliomyelitis, can be caused by other enteroviruses (34). The extremely low MS risk in

EBV-negative individuals suggests that by far most MS cases are caused by EBV and could thus potentially be prevented by a suitable vaccine. The addition of MS to the list of diseases that an EBV vaccine could target strengthens the rationale to accelerate ongoing research with the primary goal of preventing infectious mononucleosis and posttransplantation lymphoproliferative disease (35).

One of the most effective treatments for MS is anti-CD20 monoclonal antibodies, which deplete circulating memory B cells, the primary site of persistent latent EBV infection (36). This, and preliminary results obtained with EBV-specific T cell therapy (37), suggest that EBV, in addition to causing MS, contributes to MS clinical course, which could thus be potentially modified by antivirals. Directly targeting EBV could have major advantages compared with anti-CD20-based therapies, which have to be administered by intravenous infusion and may increase the risk of infections (36).

## REFERENCES AND NOTES

1. D. S. Reich, C. F. Lucchinetti, P. A. Calabresi, Multiple sclerosis. *N. Engl. J. Med.* **378**, 169–180 (2018). doi:10.1056/NEJMra1401483 Medline
2. N. L. Fransen, C.-C. Hsiao, M. van der Poel, H. J. Engelenburg, K. Verdaasdonk, M. C. J. Vincenten, E. B. M. Remmerswaal, T. Kuhlmann, M. R. J. Mason, J. Hamann, J. Smolders, I. Huitinga, Tissue-resident memory T cells invade the brain parenchyma in multiple sclerosis white matter lesions. *Brain* **143**, 1714–1730 (2020). doi:10.1093/brain/awaa117 Medline
3. D. A. Thorley-Lawson, Epstein-Barr virus: Exploiting the immune system. *Nat. Rev. Immunol.* **1**, 75–82 (2001). doi:10.1038/35095584 Medline
4. E. L. Thacker, F. Mirzaei, A. Ascherio, Infectious mononucleosis and risk for multiple sclerosis: A meta-analysis. *Ann. Neurol.* **59**, 499–503 (2006). doi:10.1002/ana.20820 Medline
5. L. I. Levin, K. L. Munger, M. V. Rubertone, C. A. Peck, E. T. Lennette, D. Spiegelman, A. Ascherio, Temporal relationship between elevation of Epstein-Barr virus antibody titers and initial onset of neurological symptoms in multiple sclerosis. *JAMA* **293**, 2496–2500 (2005). doi:10.1001/jama.293.20.2496 Medline
6. B. Serafini, B. Rosicarelli, D. Franciotta, R. Magliozi, R. Reynolds, P. Cinque, L. Andreoni, P. Trivedi, M. Salvetti, A. Faggioni, F. Aloisi, Dysregulated Epstein-Barr virus infection in the multiple sclerosis brain. *J. Exp. Med.* **204**, 2899–2912 (2007). doi:10.1084/jem.20071030 Medline
7. M. A. Moreno, N. Or-Geva, B. T. Aftab, R. Khanna, E. Croze, L. Steirman, M. H. Han, Molecular signature of Epstein-Barr virus infection in MS brain lesions. *Neuroimmunol. Neuroinflamm.* **5**, e466 (2018). doi:10.1212/NXI.0000000000000466 Medline
8. A. Hassani, J. R. Corboy, S. Al-Salam, G. Khan, Epstein-Barr virus is present in the brain of most cases of multiple sclerosis and may engage more than just B cells. *PLOS ONE* **13**, e0192109 (2018). doi:10.1371/journal.pone.0192109 Medline
9. H. Lassmann, G. Niedobitek, F. Aloisi, J. M. Middeldorp; NeuroproMiSe EBV Working Group, Epstein-Barr virus in the multiple sclerosis brain: A controversial issue—report on a focused workshop held in the Centre for Brain Research of the Medical University of Vienna, Austria. *Brain* **134**, 2772–2786 (2011). doi:10.1093/brain/awr197 Medline
10. C. L. Perdue, A. A. Eick-Cost, M. V. Rubertone, A Brief Description of the Operation of the DoD Serum Repository. *Mil. Med.* **180** (Suppl), 10–12 (2015). doi:10.7205/MILMED-D-14-00739 Medline
11. L. I. Levin, K. L. Munger, E. J. O'Reilly, K. I. Falk, A. Ascherio, Primary infection with the Epstein-Barr virus and risk of multiple sclerosis. *Ann. Neurol.* **67**, 824–830 (2010). doi:10.1002/ana.21978 Medline
12. D. H. Crawford, K. F. Macsween, C. D. Higgins, R. Thomas, K. McAulay, H. Williams, N. Harrison, S. Reid, M. Conacher, J. Douglas, A. J. Swerdlow, A cohort study among university students: Identification of risk factors for Epstein-Barr virus seroconversion and infectious mononucleosis. *Clin. Infect. Dis.* **43**, 276–282 (2006). doi:10.1086/505400 Medline
13. J. B. Dowd, A. E. Aiello, D. E. Alley, Socioeconomic disparities in the seroprevalence of cytomegalovirus infection in the US population: NHANES III. *Epidemiol. Infect.* **137**, 58–65 (2009). doi:10.1017/S0950268808000551 Medline

14. H. H. Balfour Jr., F. Sifakis, J. A. Sliman, J. A. Knight, D. O. Schmeling, W. Thomas. Age-specific prevalence of Epstein-Barr virus infection among individuals aged 6–19 years in the United States and factors affecting its acquisition. *J. Infect. Dis.* **208**, 1286–1293 (2013). doi:10.1093/infdis/jit321 Medline
15. M. Lipsitch, E. Tchetgen Tchetgen, T. Cohen, Negative controls: A tool for detecting confounding and bias in observational studies. *Epidemiology* **21**, 383–388 (2010). doi:10.1097/EDE.0b013e3181d61eeb Medline
16. V. Grut, M. Biström, J. Salzer, P. Stridh, D. Jons, R. Gustafsson, A. Fogdell-Hahn, J. Huang, N. Brenner, J. Butt, N. Bender, A. Lindam, L. Alonso-Magdalena, M. Gunnarsson, M. Vrethem, T. Bergström, O. Andersen, I. Kockum, T. Waterboer, T. Olsson, P. Sundström, Cytomegalovirus seropositivity is associated with reduced risk of multiple sclerosis—a presymptomatic case-control study. *Eur. J. Neurol.* **28**, 3072–3079 (2021). doi:10.1111/ene.14961 Medline
17. H. Tremlett, R. A. Marrie, The multiple sclerosis prodrome: Emerging evidence, challenges, and opportunities. *Mult. Scler.* **27**, 6–12 (2021). doi:10.1177/1352458520914844 Medline
18. M. Khalil, C. E. Teunissen, M. Otto, F. Piehl, M. P. Sormani, T. Gattringer, C. Barro, L. Kappos, M. Comabella, F. Fazekas, A. Petzold, K. Blennow, H. Zetterberg, J. Kuhle, Neurofilaments as biomarkers in neurological disorders. *Nat. Rev. Neurol.* **14**, 577–589 (2018). doi:10.1038/s41582-018-0058-z Medline
19. B. Delcoigne, A. Manouchehrinia, C. Barro, P. Benkert, Z. Michalak, L. Kappos, D. Leppert, J. A. Tsai, T. Plavina, B. C. Kieseier, J. Lycke, L. Alfredsson, I. Kockum, J. Kuhle, T. Olsson, F. Piehl, Blood neurofilament light levels segregate treatment effects in multiple sclerosis. *Neurology* **94**, e1201–e1212 (2020). doi:10.1212/WNL.00000000000009097 Medline
20. K. Bjornevik, K. L. Munger, M. Cortese, C. Barro, B. C. Healy, D. W. Niebuhr, A. I. Scher, J. Kuhle, A. Ascherio, Serum neurofilament light chain levels in patients with presymptomatic multiple sclerosis. *JAMA Neurol.* **77**, 58–64 (2020). doi:10.1001/jamaneurol.2019.3238 Medline
21. G. J. Xu, T. Kula, Q. Xu, M. Z. Li, S. D. Vernon, T. Ndung'u, K. Ruxrungtham, J. Sanchez, C. Brander, R. T. Chung, K. C. O'Connor, B. Walker, H. B. Larman, S. J. Elledge, Comprehensive serological profiling of human populations using a synthetic human virome. *Science* **348**, aaa0698 (2015). doi:10.1126/science.aaa0698 Medline
22. M. J. Mina, T. Kula, Y. Leng, M. Li, R. D. de Vries, M. Knip, H. Siljander, M. Rewers, D. F. Choy, M. S. Wilson, H. B. Larman, A. N. Nelson, D. E. Griffin, R. L. de Swart, S. J. Elledge, Measles virus infection diminishes preexisting antibodies that offer protection from other pathogens. *Science* **366**, 599–606 (2019). doi:10.1126/science.aay6485 Medline
23. T. J. VanderWeele, P. Ding, Sensitivity analysis in observational research: introducing the E-value. *Ann. Intern. Med.* **167**, 268–274 (2017). doi:10.7326/M16-2607 Medline
24. S. Sawcer, G. Hellenthal, M. Pirinen, C. C. Spencer, N. A. Patsopoulos, L. Moutsianas, A. Dilthey, Z. Su, C. Freeman, S. E. Hunt, S. Edkins, E. Gray, D. R. Booth, S. C. Potter, A. Goris, G. Band, A. B. Oturai, A. Strange, J. Saarela, C. Bellenguez, B. Fontaine, M. Gillman, B. Hemmer, R. Gwilliam, F. Zipp, A. Jayakumar, R. Martin, S. Leslie, S. Hawkins, E. Giannoulatou, S. D'Alfonso, H. Blackburn, F. Martinelli Boneschi, J. Liddle, H. F. Harbo, M. L. Perez, A. Spurkland, M. J. Waller, M. P. Mycko, M. Ricketts, M. Comabella, N. Hammond, I. Kockum, O. T. McCann, M. Ban, P. Whittaker, A. Kemppinen, P. Weston, C. Hawkins, S. Widaa, J. Zajicek, S. Dronov, N. Robertson, S. J. Bumpstead, L. F. Barcellos, R. Ravindrarajah, R. Abraham, L. Alfredsson, K. Ardlie, C. Aubin, A. Baker, K. Baker, S. E. Baranzini, L. Bergamaschi, R. Bergamaschi, A. Bernstein, A. Berthele, M. Boggild, J. P. Bradfield, D. Brassat, S. A. Broadley, D. Buck, H. Butzkueven, R. Capra, W. M. Carroll, P. Cavalla, E. G. Celius, S. Cepok, R. Chiavacci, F. Clerget-Darpoux, K. Clysters, G. Comi, M. Cossburn, I. Courrou-Rebeix, M. B. Cox, W. Cozen, B. A. Cree, A. H. Cross, D. Cusi, M. J. Daly, E. Davis, P. I. de Bakker, M. Debouverie, M. B. D'hooghe, K. Dixon, R. Dobosi, B. Dubois, D. Ellinghaus, I. Elovaara, F. Esposito, C. Fontenille, S. Foote, A. Franke, D. Galimberti, A. Ghezzi, J. Glessner, R. Gomez, O. Gout, C. Graham, S. F. Grant, F. R. Guerini, H. Hakonarson, P. Hall, A. Hamsten, H. P. Hartung, R. N. Heard, S. Heath, J. Hobart, M. Hoshi, C. Infante-Duarte, G. Ingram, W. Ingram, T. Islam, M. Jagodic, M. Kabesch, A. G. Kermode, T. J. Kilpatrick, C. Kim, N. Klopp, K. Koivisto, M. Larsson, M. Lathrop, J. S. Lechner-Scott, M. A. Leone, V. Leppä, U. Liljedahl, I. L. Bomfim, R. R. Lincoln, J. Link, J. Liu, A. R. Lorentzen, S. Lupoli, F. Macciardi, T. Mack, M. Marriott, V. Martinelli, D. Mason, J. L. McCauley, F. Mentch, I. L. Mero, T. Mihalova, X. Montalban, J. Motterhead, K. M. Myhr, P. Naldi, W. Ollier, A. Page, A. Palotie, J. Pelletier, L. Piccio, T. Pickersgill, F. Piehl, S. Pobylwajlo, H. L. Quach, P. P. Ramsay, M. Reunanen, R. Reynolds, J. D. Rioux, M. Rodegher, S. Roesner, J. P. Rubio, I. M. Rückert, M. Salvetti, E. Salvi, A. Santaniello, C. A. Schaefer, S. Schreiber, C. Schulze, R. J. Scott, F. Sellebjerg, K. W. Selma, D. Sexton, L. Shen, B. Simms-Acuna, S. Skidmore, P. M. Sleiman, C. Smestad, P. S. Sørensen, H. B. Søndergaard, J. Stankovich, R. C. Strange, A. M. Sulonen, E. Sundqvist, A. C. Syvänen, F. Taddeo, B. Taylor, J. M. Blackwell, P. Tienari, E. Bramon, A. Tourbah, M. A. Brown, E. Tronczynska, J. P. Casas, N. Tubridy, A. Corvin, J. Vickery, J. Jankowski, P. Villoslada, H. S. Markus, K. Wang, C. G. Mathew, J. Wason, C. N. Palmer, H. E. Wichmann, R. Plomin, E. Willoughby, A. Rautanen, J. Winkelmann, M. Wittig, R. C. Trembath, J. Yaoquanq, A. C. Viswanathan, H. Zhang, N. W. Wood, R. Zivich, P. Deloukas, C. Langford, A. Duncanson, J. R. Oksenberg, M. A. Pericak-Vance, J. L. Haines, T. Olsson, J. Hillert, A. J. Ivinson, P. L. De Jager, L. Peltonen, G. J. Stewart, D. A. Hafler, S. L. Hauser, G. McVean, P. Donnelly, A. Compston; International Multiple Sclerosis Genetics Consortium; Wellcome Trust Case Control Consortium 2, Genetic risk and a primary role for cell-mediated immune mechanisms in multiple sclerosis. *Nature* **476**, 214–219 (2011). doi:10.1038/nature10251 Medline
25. W. J. Jabs, M. Paulsen, H. J. Wagner, H. Kirchner, H. Klüter, Analysis of Epstein-Barr virus (EBV) receptor CD21 on peripheral B lymphocytes of long-term EBV-adults. *Clin. Exp. Immunol.* **116**, 468–473 (1999). doi:10.1046/j.1365-2249.1999.00912.x Medline
26. T. Olsson, L. F. Barcellos, L. Alfredsson, Interactions between genetic, lifestyle and environmental risk factors for multiple sclerosis. *Nat. Rev. Neurol.* **13**, 25–36 (2017). doi:10.1038/nrneuro.2016.187 Medline
27. H. Zdimerova, A. Murer, C. Engelmann, A. Raykova, Y. Deng, C. Gujer, J. Rühl, D. McHugh, N. Caduff, R. Naghavian, G. Pezzino, R. Capaul, A. Zbinden, G. Ferlazzo, J. D. Lünemann, R. Martin, B. Chatterjee, C. Münz, Attenuated immune control of Epstein-Barr virus in humanized mice is associated with the multiple sclerosis risk factor HLA-DR15. *Eur. J. Immunol.* (2020). Medline
28. A. Ascherio, K. L. Munger, Epidemiology of multiple sclerosis: From risk factors to prevention--An update. *Semin. Neurol.* **36**, 103–114 (2016). doi:10.1055/s-0036-1579693 Medline
29. J. M. A. Wijnands, F. Zhu, E. Kingwell, Y. Zhao, C. Evans, J. D. Fisk, R. A. Marrie, H. Tremlett, Prodrome in relapsing-remitting and primary progressive multiple sclerosis. *Eur. J. Neurol.* **26**, 1032–1036 (2019). doi:10.1111/ene.13925 Medline
30. W. A. Sibley, C. R. Bamford, K. Clark, Clinical viral infections and multiple sclerosis. *Lancet* **1**, 1313–1315 (1985). doi:10.1016/S0140-6736(85)92801-6 Medline
31. Y. Xu, K. A. Smith, A. Hiyoshi, F. Piehl, T. Olsson, S. Montgomery, Hospital-diagnosed infections before age 20 and risk of a subsequent multiple sclerosis diagnosis. *Brain* **144**, 2390–2400 (2021). doi:10.1093/brain/awab100 Medline
32. T. R. Nielsen, K. Rostgaard, N. M. Nielsen, N. Koch-Henriksen, S. Haahr, P. S. Sørensen, H. Hjalgrim, Multiple sclerosis after infectious mononucleosis. *Arch. Neurol.* **64**, 72–75 (2007). doi:10.1001/archneur.64.1.72 Medline
33. G. N. DeLorenze, K. L. Munger, E. T. Lennette, N. Orentreich, J. H. Vogelman, A. Ascherio, Epstein-Barr virus and multiple sclerosis: Evidence of association from a prospective study with long-term follow-up. *Arch. Neurol.* **63**, 839–844 (2006). doi:10.1001/archneur.63.6.noc50328 Medline
34. K. Messacar, T. L. Schreiner, J. A. Maloney, A. Wallace, J. Ludke, M. S. Oberste, W. A. Nix, C. C. Robinson, M. P. Glodé, M. J. Abzug, S. R. Dominguez, A cluster of acute flaccid paralysis and cranial nerve dysfunction temporally associated with an outbreak of enterovirus D68 in children in Colorado, USA. *Lancet* **385**, 1662–1671 (2015). doi:10.1016/S0140-6736(14)62457-Q Medline
35. J. I. Cohen, Vaccine development for Epstein-Barr virus. *Adv. Exp. Med. Biol.* **1045**, 477–493 (2018). doi:10.1007/978-981-10-7230-7\_22 Medline
36. S. L. Hauser, A. Bar-Or, G. Comi, G. Giovannoni, H.-P. Hartung, B. Hemmer, F. Lublin, X. Montalban, K. W. Rammohan, K. Selma, A. Traboulsee, J. S. Wolinsky, D. L. Arnold, G. Klingelschmitt, D. Masterman, P. Fontoura, S. Belachew, P. Chin, N. Mairon, H. Garren, L. Kappos; OPERA I and OPERA II Clinical Investigators, Ocrelizumab versus interferon beta-1a in relapsing multiple sclerosis. *N. Engl. J. Med.* **376**, 221–234 (2017). doi:10.1056/NEJMoa1601277 Medline
37. M. P. Pender, P. A. Csurhes, C. Smith, N. L. Douglas, M. A. Neller, K. K. Matthews, L. Beagley, S. Rehan, P. Crooks, T. J. Hopkins, S. Blum, K. A. Green, Z. A. Ioannides,

- A. Swayne, B. T. Aftab, K. D. Hooper, S. R. Burrows, K. M. Thompson, A. Coulthard, R. Khanna, Epstein-Barr virus-specific T cell therapy for progressive multiple sclerosis. *JCI Insight* **3**, e124714 (2018). doi:10.1172/jci.insight.124714 Medline
38. Data for: K. Bjornevik, M. Cortese, B. C. Healy, J. Kuhle, M. J. Mina, Y. Leng, S. J. Elledge, D. W. Niebuhr, A. I. Scher, K. L. Munger, A. Ascherio, Data for: K. Bjornevik, M. Cortese, B. C. Healy, J. Kuhle, M. J. Mina, Y. Leng, S. J. Elledge, D. W. Niebuhr, A. I. Scher, K. L. Munger, A. Ascherio. Longitudinal analysis reveals high prevalence of Epstein-Barr virus associated with multiple sclerosis. Harvard Dataverse (2021); <https://doi.org/10.7910/DVN/KYH9SE>.
39. U.S. Department of Defense, "2019 Demographics. profile of the military community" (DoD, 2019); <https://download.militaryonesource.mil/12038/MOS/Reports/2019-demographics-report.pdf>.
40. M. J. Knol, J. P. Vandenbroucke, P. Scott, M. Egger, What do case-control studies estimate? Survey of methods and assumptions in published case-control research. *Am. J. Epidemiol.* **168**, 1073–1081 (2008). doi:10.1093/aje/kwn217 Medline
41. A. D. Hislop, G. S. Taylor, D. Sauce, A. B. Rickinson, Cellular responses to viral infection in humans: Lessons from Epstein-Barr virus. *Annu. Rev. Immunol.* **25**, 587–617 (2007). doi:10.1146/annurev.immunol.25.022106.141553 Medline
42. S. K. Dummiere, P. S. Verghese, H. H. Balfour Jr., Primary Epstein-Barr virus infection. *J. Clin. Virol.* **102**, 84–92 (2018). doi:10.1016/j.jcv.2018.03.001 Medline
43. M. De Paschale, P. Clerici, Serological diagnosis of Epstein-Barr virus infection: Problems and solutions. *World J. Virol.* **1**, 31–43 (2012). doi:10.5501/wjv.v1.i1.31 Medline
44. J. B. Dowd, T. Palermo, J. Brite, T. W. McDade, A. Aiello, Seroprevalence of Epstein-Barr virus infection in U.S. children ages 6–19, 2003–2010. *PLOS ONE* **8**, e64921 (2013). doi:10.1371/journal.pone.0064921 Medline
45. J. M. Grimm, D. O. Schmeling, S. K. Dummiere, J. A. Knight, B. D. Mullan, J. A. Ed, R. C. Brundage, K. A. Hogquist, H. H. Balfour Jr., Prospective studies of infectious mononucleosis in university students. *Clin. Transl. Immunology* **5**, e94 (2016). doi:10.1038/cti.2016.48 Medline
46. W. Henle, G. Henle, J. Andersson, I. Ernberg, G. Klein, C. A. Horwitz, G. Marklund, L. Rymo, C. Wellinder, S. E. Straus, Antibody responses to Epstein-Barr virus-determined nuclear antigen (EBNA)-1 and EBNA-2 in acute and chronic Epstein-Barr virus infection. *Proc. Natl. Acad. Sci. U.S.A.* **84**, 570–574 (1987). doi:10.1073/pnas.84.2.570 Medline
47. G. Disanto, C. Barro, P. Benkert, Y. Naegelin, S. Schädelin, A. Giardiello, C. Zecca, K. Blennow, H. Zetterberg, D. Leppert, L. Kappos, C. Gobbi, J. Kuhle, Swiss Multiple Sclerosis Cohort Study Group, Serum Neurofilament light: A biomarker of neuronal damage in multiple sclerosis. *Ann. Neurol.* **81**, 857–870 (2017). doi:10.1002/ana.24954 Medline
48. J. Kuhle, C. Barro, U. Andreasson, T. Derfuss, R. Lindberg, Å. Sandelius, V. Liman, N. Norgren, K. Blennow, H. Zetterberg, Comparison of three analytical platforms for quantification of the neurofilament light chain in blood samples: ELISA, electrochemiluminescence immunoassay and Simoa. *Clin. Chem. Lab. Med.* **54**, 1655–1661 (2016). doi:10.1515/calm-2015-1195 Medline
49. J. Kuhle, K. Plattner, J. P. Bestwick, R. L. Lindberg, S. V. Ramagopalan, N. Norgren, A. Nissim, A. Malaspina, D. Leppert, G. Giovannoni, L. Kappos, A comparative study of CSF neurofilament light and heavy chain protein in MS. *Mult. Scler.* **19**, 1597–1603 (2013). doi:10.1177/1352458513482374 Medline
50. J. Gaiottino, N. Norgren, R. Dobson, J. Topping, A. Nissim, A. Malaspina, J. P. Bestwick, A. U. Monsch, A. Regeniter, R. L. Lindberg, L. Kappos, D. Leppert, A. Petzold, G. Giovannoni, J. Kuhle, Increased neurofilament light chain blood levels in neurodegenerative neurological diseases. *PLOS ONE* **8**, e75091 (2013). doi:10.1371/journal.pone.0075091 Medline
51. H. B. Larman, Z. Zhao, U. Laserson, M. Z. Li, A. Ciccia, M. A. M. Gakidis, G. M. Church, S. Kesari, E. M. Leproust, N. L. Solimini, S. J. Elledge, Autoantigen discovery with a synthetic human peptidome. *Nat. Biotechnol.* **29**, 535–541 (2011). doi:10.1038/nbt.1856 Medline
52. R. Vita, S. Mahajan, J. A. Overton, S. K. Dhanda, S. Martini, J. R. Cantrell, D. K. Wheeler, A. Sette, B. Peters, The Immune Epitope Database (IEDB): 2018 update. *Nucleic Acids Res.* **47** (D1), D339–D343 (2019). doi:10.1093/nar/gky1006 Medline

## ACKNOWLEDGMENTS

We thank M. Rubertone and A. Cost of the Armed Forces Health Surveillance Center, Silver Spring, MD, for control and sample identification and retrieval; N. Howard, Department of the Navy, Secretary of the Naval Council of Review Boards, Washington, DC, for MS case identification in the US Navy and Marines; D. Armitage, US Army Physical Disability Agency, Washington, DC, and L. Kent, SAIC, US Army Physical Disability Agency, Washington, DC, for MS case identification in the US Army; L. R. Hodge, US Air Force Personnel Center, San Antonio, TX, for MS case identification in the US Air Force; S. Bernstein, Uniformed Services University of the Health Sciences, Bethesda, MD, for project coordination and management; D. Leppert, University of Basel, Basel, Switzerland, for serum neurofilament measurements and interpretation of results. The views expressed are those of the authors and should not be construed to represent the positions of the Department of the Army, Department of the Navy, Department of Defense, Department of the Air Force, or Uniformed Services University of the Health Sciences. **Funding:** This work was supported by grants NS046635, NS042194, and NS103891 from the National Institute of Neurological Disorders and Stroke, National Institutes of Health (NIH); National Multiple Sclerosis Society award PP-1912-35234 and German Research Foundation (DFG) research fellowship CO 2129/1-1 to M.C.; and NIH Directors Early Independence Award DP5-OD028145 to M.J.M.. S.J.E is an investigator with the Howard Hughes Medical Institute. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. **Author contributions:** Conceptualization: K.B., M.C., K.L.M., A.A.; Data curation: D.W.N., A.I.S.; Investigation: J.K., M.J.M., Y.L., S.J.E.; Writing: K.B., M.C., K.L.M., A.A., B.C.H., J.K., M.J.M., Y.L., S.J.E., D.W.N., A.I.S. Statistical analysis: K.B., M.C., B.C.H.; Supervision: K.L.M., A.A. **Competing interests:** B.C.H. has received research support from Analysis Group, Celgene (Bristol-Myers Squibb), Verily Life Sciences, Merck-Serono, Novartis, and Genzyme. J.K. has received speaker fees, research support, travel support, and/or served on advisory boards of the Swiss MS Society, Swiss National Research Foundation (3200030\_189140/1), University of Basel, Progressive MS Alliance, Bayer, Biogen, Celgene, Merck, Novartis, Octave Bioscience, Roche, and Sanofi. M.J.M. is on the advisory board of ImmunelD and an adviser to Detect and Quantum-SI. S.J.E. is a founder of TSCAN Therapeutics, MAZE Therapeutics, Mirimus, and ImmunelD. S.J.E. serves on the scientific advisory board of Homology Medicines, TSCAN Therapeutics, MAZE, and ImmunelD, and is an adviser for MPM, none of which affect this work. S.J.E is an inventor on a patent application filed by the Brigham and Women's Hospital (US20160320406A) that covers the use of the VirScan library to identify pathogen antibodies in the blood. K.B., Y.L., A.I.S., D.W.N., A.A., K.L.M., and M.C. declare no competing interests. **Data and materials availability:** All data needed to evaluate the conclusions in this study are present in the main paper or the supplementary materials and have been deposited at the Harvard Dataverse (38).

## SUPPLEMENTARY MATERIALS

[science.org/doi/10.1126/science.abj8222](https://science.org/doi/10.1126/science.abj8222)

Materials and Methods

Figs. S1 to S4

Tables S1 and S2

References (39–52)


Data S1

MDAR Reproducibility Checklist


3 June 2021; accepted 2 December 2021

Published online 13 January 2022


10.1126/science.abj8222




**Fig. 1. Study design.** (A) Residual serum samples from the DoDSR were obtained from 810 MS patients and 1577 matched controls. We assessed whether individuals were seropositive for EBV and CMV in up to three serum samples per person. We measured sNFL in those who were EBV-negative in the first serum sample. VirScan was used to profile the virome in a subset of MS cases with serum samples collected shortly before and after symptom onset. (B) Density plot of age at onset among MS patients who were EBV-negative at time of first serum sample. The dashed line marks median age at onset. (C) Box plots of the time to first MS symptoms according to the serum sample.



**Fig. 2.** EBV infection precedes MS onset and is associated with markedly higher disease risk. (A) Proportion of individuals who were EBV-positive at the time of the first, second, and third sample. The figure is restricted to those who were EBV-negative at baseline and with EBV measurement in three samples (33 of 35 MS patients and 90 of 107 controls). A significantly higher proportion of individuals who later developed MS were EBV-positive in the second (28 of 33 MS patients) and third (32 of 33 MS patients) sample compared with individuals who did not develop MS (second sample: 40 of 90 controls; third sample: 51 of 90 controls). \*\*\* $P < 0.0001$ , two-sided Fisher's exact test. (B) Proportion of individuals who were CMV-positive at the time of the first, second, and third sample collected in the study. The figure is restricted to those who were CMV- and EBV-negative at baseline. The proportion who were CMV-positive was similar in the second (two of 23 MS patients versus four of 60 controls) and third sample (three of 23 MS patients versus seven of 60 controls). All  $P > 0.05$ , two-sided Fisher's exact test. (C) Risk ratio for MS according to EBV status. EBV seroconversion by the time of the third sample and EBV seropositivity at the time of the first sample were associated with a 32-fold and 26-fold increased risk of developing MS, respectively, in matched analyses. \*\* $P < 0.01$  and \*\*\* $P < 0.001$ , two-sided univariable conditional logistic regression model. (D) Risk ratio for MS according to CMV status. \*\* $P < 0.01$ , two-sided univariable conditional logistic regression model.



**Fig. 3. EBV infection precedes elevation of sNfL before the onset of MS.** (A) Box plots of sNfL levels before, around, and after the time of EBV infection. \* $P < 0.05$ , two-sided multivariable linear regression model adjusted for age and sex. (B) Within-person increase in sNfL levels in MS cases around and after time of EBV infection compared with before EBV infection. \*\* $P < 0.01$ , two-sided linear mixed-effects regression model. (C) Within-person increase in sNfL levels in controls around and after time of EBV infection compared with before EBV infection. Error bars in (B) and (C) are 95% CIs. sNfL levels increased significantly more in MS cases than in controls in the sample collected after time of EBV infection compared with before EBV infection ( $P < 0.001$ , two-sided linear mixed-effects regression model).



**Fig. 4. Antibodies against EBV peptides in MS cases detected using virome-wide screening.** (A to D) Scatter plots showing the mean Z scores of serum antibody enrichment against the entire viral peptidome in cases and controls in the pre- and post-onset samples. Each point represents one peptide. (E and F) Number of viral peptides with nominally significantly different antibody binding between cases and controls in the pre- and post-onset samples (using a Z score  $>3.5$  as an enrichment cut-off; compared using two-sided Fisher's exact test) and their mapping to viral species. (G and H) Mapping of the statistically significant EBV peptides to EBV antigens (for details, see tables S1 and S2).

## Longitudinal analysis reveals high prevalence of Epstein-Barr virus associated with multiple sclerosis

Kjetil Bjornevik<sup>1</sup>Marianna Cortese<sup>2</sup>Brian C. Healy<sup>3</sup>Jens Kuhle<sup>4</sup>Michael J. Mina<sup>5</sup>Yumei Leng<sup>6</sup>Stephen J. Elledge<sup>7</sup>David W. Niebuhr<sup>8</sup>Ann I. Scher<sup>9</sup>Kassandra L. Munger<sup>10</sup>Alberto Ascherio<sup>11</sup>

Science, **Ahead of Print**

**View the article online**

<https://www.science.org/doi/10.1126/science.abj8222>

**Permissions**

<https://www.science.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of service](#)